

Cerebrovascular Response to CO₂ Following 10 Days of Intermittent Hypoxia in Humans

Jordan S. Querido; Joseph F. Welch; Najib T. Ayas; A. William Sheel

- INTRODUCTION:** It has been demonstrated that the cerebrovascular response to hypoxia is blunted following 10 d of intermittent hypoxia (IH) in healthy humans. The purpose of this study was to test the hypothesis that IH reduces the cerebrovascular response to CO₂.
- METHODS:** Healthy male subjects ($N = 8$; 25 ± 2 yr) were exposed to 10 consecutive days of IH (12% O₂ for 5 min followed by 5 min of normoxia for 1 h/d). The cerebrovascular response to CO₂ was assessed prior to (PRE-IH) and following (POST-IH) the IH paradigm with transcranial Doppler ultrasound.
- RESULTS:** There was no change in eupnic measures during or following the IH paradigm; however, the ventilatory response to IH increased by the last exposure (3.0 ± 2.8 L · min⁻¹). Cerebral blood flow velocity decreased and increased with hypocapnia and hypercapnia, respectively, but cerebrovascular sensitivity to CO₂ remained unchanged with IH (PRE-IH: $2.58 \pm 0.50\%/mmHg$; POST-IH: $2.59 \pm 0.74\%/mmHg$).
- DISCUSSION:** Our data indicates that 10 d of IH in healthy humans does not alter the cerebrovascular response to CO₂. Redundancy of cerebrovascular regulation mechanisms to CO₂ may work to counteract IH-induced dysregulation and protect cerebral tissue.
- KEYWORDS:** hypoxemia, cerebrovascular control, carbon dioxide.

Querido JS, Welch JF, Ayas NT, Sheel AW. Cerebrovascular response to CO₂ following 10 days of intermittent hypoxia in humans. *Aerosp Med Hum Perform*. 2015; 86(9):782–786.

Exposure to intermittent hypoxia (IH) is considered to be unique from sustained hypoxia, whereby the inability to adapt to IH often leads to autonomic dysregulation.¹⁸ For instance, isolated rodent vessels exhibit autonomic vasodilator dysregulation following IH.¹⁹ Similarly, patients with obstructive sleep apnea demonstrate autonomic dysfunction in the ventilatory, cardiovascular, and cerebrovascular systems—an effect attributed to the IH inherent to the syndrome.⁵ Healthy human models of IH show autonomic dysregulation that parallels pathology.^{16,22} In a previous study, we found the cerebrovascular sensitivity to acute hypoxia was significantly blunted following 10 consecutive days of IH in healthy humans.²² The mechanistic explanation for this impairment in hypoxic cerebrovascular regulation is still uncertain, but may be the result of an increase in IH-induced oxidative stress, leading to endothelial dysfunction.¹³ Oxidative stress due to IH may increase reactive oxygen species, which in turn reduces the bioavailability of endothelial nitric oxide (NO)—a key regulator of cerebrovasculature in hypoxia.¹³ Support for this hypothesis is demonstrated in studies that have prevented the typical

IH-induced autonomic impairments with the administration of an antioxidant.^{7,14}

Although multifactorial, cerebrovascular regulation is particularly sensitive to adjustments in arterial pressures of CO₂.⁸ Similar to the cerebrovascular response to hypoxia, endothelial function plays a significant role in the cerebrovascular response to hypercapnia via a CO₂-NO pathway.^{12,26} However, the effect of IH on the cerebrovascular response to CO₂ has shown conflicting results, being increased,¹¹ decreased,²⁰ or unchanged with IH.¹⁰ Integrating and interpreting the available literature is difficult, given that differences in IH protocol (e.g., short- vs. long-duration hypoxia cycles) and subject characteristics

From the Faculty of Medicine and the School of Kinesiology, University of British Columbia–Vancouver, Vancouver, British Columbia, Canada.

This manuscript was received for review in October 2014. It was accepted for publication in June 2015.

Address correspondence to: A. William Sheel, 6108 Thunderbird Blvd., Vancouver, BC, Canada V6T 1Z3; bill.sheel@ubc.ca.

Reprint & Copyright © by the Aerospace Medical Association, Alexandria, VA.

DOI: 10.3357/AMHP.4192.2015

(e.g., healthy humans vs. patients) could play a large role in the discrepancies between studies. Accordingly, the purpose of this study was to investigate cerebrovascular regulation to CO₂ following an IH protocol that we previously demonstrated elicited cerebral blood flow dysregulation.²²

Our healthy human model of IH led to cerebrovascular dysregulation in acute hypoxia, a likely result of endothelial dysfunction from a reduced bioavailability of NO.²² If NO plays a key role in the cerebrovascular response to CO₂, then it is expected that this identical IH paradigm would result in a blunted cerebral blood flow response to CO₂. Therefore, it is hypothesized that the cerebrovascular sensitivity to CO₂ is reduced following 10 consecutive days of IH in healthy humans.

METHODS

Subjects

Healthy young male subjects ($N = 8$; 25 ± 2 yr) were recruited to participate in the study after providing written informed consent. All subjects (183 ± 6 cm; 77 ± 10 kg) were free of any known cardiorespiratory illness. Subjects underwent a preliminary session to familiarize themselves with the experimental setup and procedures. All procedures and protocols were approved by the Clinical Research Ethics Board at the University of British Columbia, which conforms to the Declaration of Helsinki.

Equipment

Breathing frequency, tidal volume, and minute ventilation (V_I) were determined from a pneumotachograph (model 3813, Hans Rudolph, Kansas City, MO). Ventilated O₂ and CO₂ were measured at the mouth and analyzed using gas analyzers (models S-3A/I and CD-A, respectively, Applied Electrochemistry, Pittsburgh, PA). Beat-by-beat blood pressure (including mean arterial pressure) and oxyhemoglobin saturation (S_pO_2) were measured noninvasively at the finger with photoplethysmography (Finometer, FMS, Arnhem, Netherlands) and finger pulse oximetry (Model 3740, Ohmeda, Louisville, CO), respectively. Standard 3-lead electrocardiography was used to determine heart rate. Cerebral blood flow velocity was continuously measured at the proximal segment of the middle cerebral artery with a 2 MHz pulsed-wave transcranial Doppler ultrasound (Neurovision 500 M, Multigon Industries, Yonkers, NY). All data was collected using an analog to digital converter (PowerLab/16SP ML 795, ADInstruments, Colorado Springs, CO), sampled at 200 Hz and stored on a computer for offline analysis (Chart V5.02, ADInstruments).

Procedure

The cerebrovascular sensitivity to CO₂ was assessed on the day prior to (PRE-IH) and following (POST-IH) 10 consecutive days of IH. The middle cerebral artery was insonated through the right temporal window, above the zygomatic arch, to obtain backscattered Doppler signals. Optimal Doppler signals were achieved using previously described methods.²¹ A transparency of each subject's facial features was taken

to ensure consistency of probe placement during the two experimental sessions (PRE-IH and POST-IH). The probe was then secured using a headband device (Marc 600, Spencer Technologies, Seattle, WA) to provide a fixed angle of insonation. For each subject, the same depth and gain of the Doppler signal was used during PRE-IH and POST-IH testing. Placement of the Doppler probe was performed by the same investigator. Each experimental day began with a minimum of 10 min of resting eupnea to ensure stable baseline measurements. Subjects abstained from caffeine, alcohol, and exhaustive exercise for 24 h prior to the experimental tests. The IH protocol consisted of subjects breathing a hypoxic inspirate [fraction of inspired oxygen (F_{IO_2}) of 12%, corresponding to an approximate altitude of 13,123 ft (4000 m)] from a reservoir for 5 min followed by 5 min of room-air breathing; this cycle was repeated for 1 h each day (Fig. 1). Subjects wore a sealed facemask during hypoxic exposures and breathed through a mouthpiece during sensitivity




Fig. 1. Physiological measurements during an IH exposure in a representative subject.

testing. Ventilated gases were continuously monitored at the mouth to ensure there were no leaks in the inspiratory circuit and precise hypoxic control was attained. The cerebrovascular sensitivity to CO₂ was determined by the rebreathe method as described by Duffin *et al.*³ Subjects were required to hyperventilate for 5 min in order to reduce end-tidal partial pressure of CO₂ (P_{ET}CO₂) by ~15 mmHg. Following the hyperventilation period, subjects expired completely before taking three deep breaths from the rebreathing bag, consisting of 7% CO₂ and balance O₂. Subjects then continued to rebreathe from the bag at a self-paced rate. During the rebreathe portion of the test, O₂ pressure of the inspire was maintained at 150 mmHg by a computer-controlled device. The test was terminated once P_{ET}CO₂ reached 60 mmHg.

Statistical Analysis

For each measured physiological variable, an average was determined at each P_{ET}CO₂ in 1-mmHg intervals. The slope of the linear regression between each variable and P_{ET}CO₂ was taken to represent the sensitivity to CO₂ (Fig. 2). For ventilatory measures, linear regression began at the ventilatory threshold. Student's paired *t*-tests were used to investigate the differences in physiological variables during eupnea between PRE-IH and POST-IH experimental sessions, as well as the differences in sensitivities to CO₂ (Statistica V7, Statsoft Inc., Tulsa, OK). Pearson product moment correlations were performed on selected dependent variables. The level of significance was set at *P* < 0.05 for all statistical comparisons. All data is presented as mean ± SD.

RESULTS

All subjects completed the entire experimental protocol. There was no effect of IH on eupnic measures during the 10-d IH protocol. Within an IH exposure, the hypoxic inspire was well-controlled (mean F_IO₂ = 12.15 ± 0.3%, corresponding to a




Fig. 2. Cerebrovascular responsiveness to CO₂ PRE-IH (black circles) and POST-IH (white circles) in a representative subject. MCA_{Vmean} = mean middle cerebral artery velocity.

P_aO₂ of 87 mmHg), resulting in a decrease and increase in S_pO₂ and V_I, respectively (Fig. 1).

Specifically, on the final day of IH exposure, the hypoxia resulted in a S_pO₂ of 84.7 ± 2.6% and a subsequent V_I of 15.6 ± 3.0 L · min⁻¹. The V_I response to hypoxia significantly increased from the first to last day of IH exposures [3.0 ± 2.8 L · min⁻¹ increase; *t*(7) = -2.47, *P* = 0.046]. There was no change in any physiological measure during eupnea between PRE-IH and POST-IH experimental sessions (Table I).

The cerebrovascular sensitivity to CO₂ was 2.58 ± 0.50%/mmHg PRE-IH and did not change with IH [POST-IH: 2.59 ± 0.74%/mmHg; *t*(7) = -0.03, *P* = 0.97; Fig. 3]. Similarly, there was no effect of IH on the CO₂ sensitivity of any other ventilatory or cardiovascular measures (*P* > 0.05).

DISCUSSION

We assessed the cerebrovascular response to CO₂ in healthy humans following 10 consecutive days of IH. Contrary to our previous report that demonstrated a reduced cerebrovascular response to acute hypoxia following an identical IH paradigm, cerebrovascular reactivity to CO₂ was unchanged, although there were adjustments in the ventilatory response to IH.

IH is a stimulus that leads to multiple adjustments in ventilatory, cardiovascular, neural, and cerebrovascular control.^{1,6} Animal, healthy human, and patient models of IH have demonstrated blunted or augmented autonomic responsiveness depending on the autonomic system being investigated.^{5,19,22} Although the physiological effects of IH are well-described, the available literature does show some inconsistencies. For example, improvements and impairments in cerebrovascular reactivity have been shown following IH,^{11,22} leading some to suggest a therapeutic effect of IH rather than pathological.²⁴ A central caveat to integrating the results of the literature is the disparity between IH paradigms employed, an important consideration given that a distinct IH paradigm mediates the physiological response.¹⁸ In particular, the severity of the hypoxic exposures and the number of hypoxic iterations (i.e., hypoxia-reoxygenation cycles) appear to be important in eliciting a pathological rather than therapeutic response.¹⁸ The present study used an IH paradigm previously demonstrated

Table I. Physiological Measurements During Eupnea in the PRE-IH and POST-IH Experimental Sessions.

	PRE-IH	POST-IH
Fb (breath/min)	13.9 ± 4.4	13.6 ± 5.5
V _T (L)	0.94 ± 0.21	0.97 ± 0.24
V _I (L · min ⁻¹)	12.4 ± 3.4	12.3 ± 3.4
P _{ET} CO ₂ (mmHg)	39 ± 2	38 ± 2
HR (bpm)	59 ± 8	59 ± 8
MAP (mmHg)	92 ± 5	94 ± 5
MCA _{Vmean} (cm · s ⁻¹)	62 ± 9	60 ± 11

IH = intermittent hypoxia; Fb = breathing frequency; V_T = tidal volume; V_I = inspired minute ventilation; P_{ET}CO₂ = end-tidal partial pressure of CO₂; HR = heart rate; MAP = mean arterial pressure; MCA_{Vmean} = mean blood flow velocity of the middle cerebral artery.




Fig. 3. The effect of IH on cerebrovascular responsiveness to CO₂. Error bars represent SD. MCAV_{mean} = mean middle cerebral artery velocity.

to cause cerebrovascular dysfunction in acute hypoxia. However, our results demonstrated no effect of IH on cerebrovascular response to CO₂, suggesting that diverse mechanisms mediate cerebrovascular control in hypoxia and CO₂.

The cerebrovasculature is particularly sensitive to changes in arterial pressure of CO₂, with increases and decreases leading to cerebral vessel dilation and constriction, respectively.⁸ Previous investigators have demonstrated the importance of NO in mediating the cerebrovascular response to CO₂.²⁶ Given the reduced bioavailability of NO following IH,⁴ we expected a blunted cerebrovascular response to CO₂ following our IH paradigm. Surprisingly, our data shows no effect of IH on cerebrovascular responsiveness to CO₂; however, we are not the first to show this.^{10,27} An explanation for the absence of an effect of IH is uncertain, but the multifaceted and redundant nature of cerebrovascular control may play a role. For instance, it is theorized that CO₂ directly influences cerebrovascular tone independently of any intermediates (e.g., NO).²³ It is possible that an impairment in cerebrovascular chemoregulation via NO unavailability could have been compensated for by a direct CO₂ influence on the cerebral vessels. In this sense, previous studies that demonstrated a blunted cerebrovascular responsiveness to CO₂ in patients with obstructive sleep apnea may be explained by a desensitization of cerebral CO₂ chemosensors from the apnea-induced periods of hypercapnia.^{2,20} Although the poikilocapnic IH in the current study may have reduced the bioavailability of NO, a direct local effect of CO₂ on cerebral vessels could have maintained proper autonomic function, thereby negating the influence of NO on cerebrovascular response to hypoxia.⁹ This implies a critical role of CO₂ on the physiological consequences of IH. Data from breath-hold divers, on the other hand, have shown normal cerebrovascular responsiveness to CO₂,¹⁰ thus, an isolated intermittent hypo-/hypercapnia paradigm may provide further insights into the mechanisms of cerebrovascular control to CO₂.

There are inherent limitations in the current study. Transcranial Doppler ultrasonography provides an estimation of

cerebral blood flow with the assumption that the insonated vessel maintains a fixed diameter. Previous reports have demonstrated no change in cross-sectional area of the middle cerebral artery during CO₂ challenges.^{21,25} As a result, we consider our transcranial Doppler measurement to represent real changes in cerebral blood flow due to cerebral vessel dilation and constriction of small resistance vessels downstream of the area of insonation. In addition, extending our results to other models of IH must be done with caution. As previously mentioned, the specific IH paradigm combined with confounding factors can mediate the physiological response. While not the focus of the current study, the effect of prior IH exposure on subjective measures of performance and cognition in acute hypoxia has been the topic of previous investigations.^{15,17} Further research investigating this possible effect could prove useful for practically relevant models of hypoxia, such as aviation and high-altitude.

In summary, we previously demonstrated cerebrovascular dysregulation in acute hypoxia following 10 d of IH. In the current study, we used an identical IH paradigm and found no effect on the cerebrovascular response to CO₂, although ventilatory responsiveness to hypoxia increased. Our data suggest a possible redundancy in cerebrovascular control during CO₂ challenges, which may act to protect the cerebrovasculature from IH-induced impairments.

ACKNOWLEDGMENTS

This study was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and a Canadian Institutes of Health Research (CIHR) Respiratory Sleep Disorders Research Team Grant. J. F. Welch was supported by a Four Year Fellowship from the University of British Columbia. A. W. Sheel was supported by a New Investigator Award from the CIHR.

The authors declare that they have no conflict of interest.

Authors and affiliations: Jordan S. Querido, Ph.D., and Najib T. Ayas, M.D., Faculty of Medicine, and Joseph F. Welch, M.Res., and A. William Sheel, Ph.D., School of Kinesiology, University of British Columbia-Vancouver, Vancouver, British Columbia, Canada.

REFERENCES

1. Casas M, Casas H, Pages T, Rama R, Ricart A, et al. Intermittent hypobaric hypoxia induces altitude acclimation and improves the lactate threshold. *Aviat Space Environ Med.* 2000; 71(2):125–130.
2. Diomedes M, Placidi F, Cupini LM, Bernardi G, Silvestrini M. Cerebral hemodynamic changes in sleep apnea syndrome and effect of continuous positive airway pressure treatment. *Neurology.* 1998; 51(4): 1051–1056.
3. Duffin J, Mohan RM, Vasiliou P, Stephenson R, Mahamed S. A model of the chemoreflex control of breathing in humans: model parameters measurement. *Respir Physiol.* 2000; 120(1):13–26.
4. Feng J, Zhang D, Chen B. Endothelial mechanisms of endothelial dysfunction in patients with obstructive sleep apnea. *Sleep Breath.* 2012; 16(2):283–294.
5. Foster GE, Hanly PJ, Ostrowski M, Poulin MJ. Effects of continuous positive airway pressure on cerebral vascular response to hypoxia in patients with obstructive sleep apnea. *Am J Respir Crit Care Med.* 2007; 175(7):720–725.

6. Foster GE, Poulin MJ, Hanly PJ. Intermittent hypoxia and vascular function: implications for obstructive sleep apnoea. *Exp Physiol.* 2007; 92(1):51–65.
7. Grebe M, Eisele HJ, Weissmann N, Schaefer C, Tillmanns H, et al. Antioxidant vitamin C improves endothelial function in obstructive sleep apnea. *Am J Respir Crit Care Med.* 2006; 173(8):897–901.
8. Heistad DD, Kontos HA. Cerebral circulation. In: Shepherd JT, Abboud FM, Geiger SR, editors. *Handbook of physiology. The cardiovascular system. Peripheral circulation. III.* Bethesda (MD): American Physiological Society; 1983:137–182.
9. Ide K, Worthley M, Anderson T, Poulin MJ. Effects of the nitric oxide synthase inhibitor L-NMMA on cerebrovascular and cardiovascular responses to hypoxia and hypercapnia in humans. *J Physiol.* 2007; 584(Pt. 1):321–322.
10. Ivancev V, Palada I, Valic Z, Obad A, Bakovic D, et al. Cerebrovascular reactivity to hypercapnia is unimpaired in breath-hold divers. *J Physiol.* 2007; 582(Pt. 2):723–730.
11. Kolb JC, Ainslie PN, Ide K, Poulin MJ. Effects of five consecutive nocturnal hypoxic exposures on the cerebrovascular responses to acute hypoxia and hypercapnia in humans. *J Appl Physiol.* 2004; 96(5): 1745–1754.
12. Lavi S, Gaitini D, Milloul V, Jacob G. Impaired cerebral CO₂ vasoreactivity: association with endothelial dysfunction. *Am J Physiol Heart Circ Physiol.* 2006; 291(4):H1856–H1861.
13. Lavie L. Sleep-disordered breathing and cerebrovascular disease: a mechanistic approach. *Neurol Clin.* 2005; 23(4):1059–1075.
14. Lee DS, Badr MS, Mateika JH. Progressive augmentation and ventilatory long-term facilitation are enhanced in sleep apnoea patients and are mitigated by antioxidant administration. *J Physiol.* 2009; 587(Pt. 22): 5451–5467.
15. Leiffen D, Poquin D, Savourey G, Barraud PA, Raphel C, Bittel J. Cognitive performance during short acclimation to severe hypoxia. *Aviat Space Environ Med.* 1997; 68(11):993–997.
16. Lusina SJ, Kennedy PM, Inglis JT, McKenzie DC, Ayas NT, Sheel AW. Long-term intermittent hypoxia increases sympathetic activity and chemosensitivity during acute hypoxia in humans. *J Physiol.* 2006; 575(Pt. 3):961–970.
17. MacNutt MJ, Laursen PB, Kedia S, Neupane M, Parajuli P, et al. Acclimatisation in trekkers with and without recent exposure to high altitude. *Eur J Appl Physiol.* 2012; 112(9):3287–3294.
18. Navarrete-Opazo A, Mitchell GS. Therapeutic potential of intermittent hypoxia: a matter of dose. *Am J Physiol Regul Integr Comp Physiol.* 2014; 307(10):R1181–R1197.
19. Phillips SA, Olson EB, Morgan BJ, Lombard JH. Chronic intermittent hypoxia impairs endothelium-dependent dilation in rat cerebral and skeletal muscle resistance arteries. *Am J Physiol Heart Circ Physiol.* 2004; 286(1):H388–H393.
20. Placidi F, Diomedes M, Cupini LM, Bernardi G, Silvestrini M. Impairment of daytime cerebrovascular reactivity in patients with obstructive sleep apnoea syndrome. *J Sleep Res.* 1998; 7(4):288–292.
21. Poulin MJ, Robbins PA. Indexes of flow and cross-sectional area of the middle cerebral artery using Doppler ultrasound during hypoxia and hypercapnia in humans. *Stroke.* 1996; 27(12):2244–2250.
22. Querido JS, Godwin JB, Sheel AW. Intermittent hypoxia reduces cerebrovascular sensitivity to isocapnic hypoxia in humans. *Respir Physiol Neurobiol.* 2008; 161(1):1–9.
23. Ringelstein EB, Otis CL. Physiological testing of vasomotor reserve. In: Newell DW, Aaslid R, editors. *Transcranial Doppler.* New York: Raven Press; 1992.
24. Serebrovskaya TV, Manukhina EB, Smith ML, Downey HF, Mallet RT. Intermittent hypoxia: cause of or therapy for systemic hypertension? *Exp Biol Med (Maywood).* 2008; 233(6):627–650.
25. Serrador JM, Picot PA, Rutt BK, Shoemaker JK, Bondar RL. MRI measures of middle cerebral artery diameter in conscious humans during simulated orthostasis. *Stroke.* 2000; 31(7):1672–1678.
26. Thompson BG, Pluta RM, Gorton ME, Oldfield EH. Nitric oxide mediation of chemoregulation but not autoregulation of cerebral blood flow in primates. *J Neurosurg.* 1996; 84(1):71–78.
27. Urbano F, Roux F, Schindler J, Mohsenin V. Impaired cerebral autoregulation in obstructive sleep apnea. *J Appl Physiol.* 2008; 105(6):1852–1857.