

Action Boundary Proximity Effects on Perceptual-Motor Judgments

Caleb D. Johnson; Alice D. LaGoy; Gert-Jan Pepping; Shawn R. Eagle; Anne Z. Beethe; Joanne L. Bower; Candice A. Alfano; Richard J. Simpson; Christopher Connaboy

- INTRODUCTION:** Designed as a more ecological measure of reaction times, the Perception-Action Coupling Task (PACT) has shown good reliability and within-subject stability. However, a lengthy testing period was required. Perceptual-motor judgments are known to be affected by proximity of the stimulus to the action boundary. The current study sought to determine the effects of action boundary proximity on PACT performance, and whether redundant levels of stimuli, eliciting similar responses, can be eliminated to shorten the PACT.
- METHODS:** There were 9 men and 7 women who completed 4 testing sessions, consisting of 3 familiarization cycles and 6 testing cycles of the PACT. For the PACT, subjects made judgments on whether a series of balls presented on a tablet afford “posting” (can fit) through a series of apertures. There were 8 ratios of ball to aperture size (B-AR) presented, ranging from 0.2 to 1.8, with each ratio appearing 12 times (12 trials) per cycle. Reaction times and judgment accuracy were calculated, and averaged across all B-ARs. Ratios and individual trials within each B-AR were systematically eliminated. Variables were re-averaged, and intraclass correlation coefficients (ICC) and coefficients of variation (CV_{TE}) were calculated in an iterative manner.
- RESULTS:** With elimination of the 0.2 and 1.8 B-ARs, the PACT showed good reliability ($ICC = 0.81-0.99$) and consistent within-subject stability ($CV_{TE} = 2.2-14.7\%$). Reliability ($ICC = 0.81-0.97$) and stability ($CV_{TE} = 2.6-15.6\%$) were unaffected with elimination of up to 8 trials from each B-AR.
- DISCUSSION:** The shortened PACT resulted in an almost 50% reduction in total familiarization/testing time required, significantly increasing usability.
- KEYWORDS:** PACT, Perception-Action Coupling Task, affordance, reliability, reaction time.

Johnson CD, LaGoy AD, Pepping G-J, Eagle SR, Beethe AZ, Bower JL, Alfano CA, Simpson RJ, Connaboy C. Action boundary proximity effects on perceptual-motor judgments. *Aerospace Med Hum Perform*. 2019; 90(12):1000-1008.

An action boundary can be defined as a critical point that limits a particular action. That is, an action boundary is the point where an individual's capabilities are exceeded by the movement demands and the actor must adjust their movement strategy for the resulting action to be successful.¹⁰ Action boundaries are derived from Gibson's theory of affordances, or the possibilities for action within a given environment.¹¹ For example, the action boundary for an individual deciding whether they can step up on a raised surface would be the maximum surface height for which they could successfully push themselves up onto using their legs. If they encountered a surface that was above this height, they would be forced to select another movement strategy for successful action, such as pushing themselves up on the surface with their arms.

The ability of an individual to accurately judge their action boundaries and alter their behavior accordingly reflects their

accuracy of perceptual-motor control and has broad implications for the successful control of movement, agency, and decision making.²⁴ Inadequate attunement to these boundaries, defined as the process of calibrating an individual to their action boundaries, has been shown to result in altered postural control and movement patterns, increased latency in reaction time measures, and overall decreases in task performance.^{8,17,18} Summarily, the level to which someone is accurately attuned to

From the Neuromuscular Research Laboratory, Department of Sports Medicine and Nutrition, University of Pittsburgh, Pittsburgh, PA, USA.

This manuscript was received for review in March 2019. It was accepted for publication in September 2019.

Address correspondence to: Christopher Connaboy, Ph.D., 3860 S. Water St., Pittsburgh, PA, 15217; connaboy@pitt.edu.

Reprint & Copyright © by the Aerospace Medical Association, Alexandria, VA.

DOI: <https://doi.org/10.3357/AMHP.5376.2019>

their action boundaries likely holds significant implications for limiting behavioral risk, relating to their ability to recognize and avoid risky movement behaviors. Military personnel, including pilots and astronauts, must continually integrate sources of perceptual information and make movement judgments that have significant implications in terms of behavioral risk.^{1,4} Further, these judgments often must be made under extreme environmental or operational conditions.^{1,4,23} Therefore, perceptual attunement to action boundaries would seem to hold high applicability to these populations.

However, a lack of literature connecting performance-based outcomes to psychological or behavioral health is a critical gap in research on affordances and behavioral risk. Traditional performance measures for alertness or reaction time fail to incorporate any form of perceptual-motor judgment, potentially accounting for this gap in literature.^{5,15,16} While traditional measures require quick responses, they fail to incorporate the critical notions of action boundaries or spatial judgments. Their decisional components are either nonexistent (i.e., simply moving a finger/mouse when the stimulus is presented) or limited to binary choices (i.e., go / no-go tasks).^{5,15,16}

The Perception-Action Coupling Task (PACT) software was designed as an ecologically valid measure of decision-making, with the inclusion of action boundary judgments as the reactionary stimuli.^{7,9,21} Based on the task first described by Smith and Pepping,²¹ the PACT is performed on a touchscreen tablet and requires individuals to judge whether a virtual ball can fit inside a virtual hole (aperture). During the assessment, eight ratios of ball-to-aperture size are presented ranging from 0.2 to 1.8 (**Fig. 1**). The action boundary is represented by a ball-to-aperture ratio (B-AR) of 1.0, where the action of moving the ball into the hole is just afforded. At any ratio higher than 1.0, the action would no longer be possible (unafforded). The initial work done to establish the intersession reliability and within-subject stability associated with the PACT is described in detail elsewhere.⁷ The PACT was shown to be reliable and stable over repeated administrations; however, it required relatively lengthy familiarization (15 min) and data collection (5–10 min) periods which may limit usability and induce testing fatigue.⁷

Previous literature has demonstrated more consistency in performance of action boundary accuracy assessments when an individual is asked to make judgments with respect to stimuli that are far from a given action boundary. However, as the stimuli approach the action boundary, increased variability and decrements in performance are observed.^{18,21,22} Subsequently, the B-ARs that are furthest from the action boundary on the PACT (i.e., 0.2, 0.4, 1.6, and 1.8) likely have the lowest ability to discern between individuals who perform better (quicker and more accurate) and worse on the task. Further, they may elicit redundant responses with the ratio closest to them (i.e., 0.2 with 0.4), given that they present the easiest perceptual judgments. In other words, on average, participants may perform similarly in response to B-ARs of 0.4 and 0.2, even though 0.4 is technically slightly closer to the action boundary.

Given the lengthy time required for familiarization and testing with the PACT, the aim of the current study was to explore ways to reduce the necessary testing time of the PACT thereby increasing its usability. In this effort, two experiments were conducted. Based on the relationship between response times and accuracy and the action boundary described above, the purpose of Experiment 1 was to determine the effects of eliminating B-ARs on the intersession reliability and within-subject stability of the PACT. Data from a previous longitudinal study was recalculated for analysis after iteratively eliminating several B-ARs. We hypothesized that several ratios furthest from the action boundary could be eliminated without affecting the representative behaviors on the PACT described by previous work.⁷ Further, we hypothesized that eliminating these ratios would not change the inherent continuum of reaction times and accuracy of judgments about the action boundary.

EXPERIMENT 1

METHODS

Subjects

The complete methods and experimental design for the current study are described in detail elsewhere.⁷ Briefly, 16 subjects (9 men, 7 women; age = 27.8 ± 3.6 yr) were recruited with the inclusion criteria of: ages 18–40, having corrected 20/20 vision, free from any visual impairments, and not taking medications that would impair cognitive processes, alertness or vision. Subjects completed 4 testing sessions, separated by at least 6 d (Mean = 9.67 ± 3.4 d). The study protocol was approved in advance by the University of Pittsburgh Institutional Review Board. Each subject provided written informed consent before participating.

Equipment and Procedures

The PACT was performed on a touchscreen tablet (iPad 3rd Gen.; Apple, Cupertino, CA). Each testing session was identical and consisted of the completion of 9 cycles of the PACT, in a quiet environment with minimal distractions. To perform the PACT (**Fig. 1**), subjects begin with their index or middle finger on the start button. At a randomized interval (0.01–0.70 s), a virtual ball and aperture appear on the screen. On presentation of the ball-aperture pairing, subjects make their judgment on whether the ball affords posting (can fit) through the aperture, move their finger to the joystick, and either swipe forward if they judge that the ball can fit through the aperture, or backward if they judge it cannot. Subjects are not given any feedback as to the accuracy of their judgments. Across each cycle of the PACT, each B-AR is presented 16 times, in a randomized order. Each cycle lasts approximately 5 min, depending on the speed of responses.

Several variables are calculated from responses on the PACT. Accuracy of judgments (ACC) is calculated as the percentage of total trials where an accurate judgment was made, with an accurate judgment defined as one where either the ball afforded posting and the subject moved the joystick




Fig. 1. Depiction of PACT interface and example ball-to-aperture ratios. Ball = gray circle at bottom of screen; aperture = white circle at top of screen; start button = circle with fingerprint on bottom right; joystick = circle to left of start button. Grayscale is used for the print version; the normal PACT interface is colored with a yellow ball, white aperture, green start button, and blue joystick.

forward, or the ball did not afford posting and the subject moved the joystick backward. Reaction time (RT) was determined by the time between presentation of the B-AR and the subject making a movement from the start button. Movement time (MT) was determined by the time between the subject making a movement from the start button and initiating a movement with the joystick. Initiation time (IT) was determined by the time between initiation and completion of a movement with the joystick. Finally, response time (RST) was determined by the sum of the previous three variables: RT, MT and IT.

As mentioned previously (Sec. 1.0), the study by Connaboy *et al.*⁷ found that three cycles were required for familiarization at the beginning of each session, and one to two further cycles of testing were necessary to obtain a stable, reliable measure across all variables. Therefore, the first three cycles of testing were eliminated, and all variables were calculated over the 4th cycle (1 cycle of testing, 128 trials) and the 4th and 5th cycles combined (2 cycles of testing, 256 trials). Finally, variables were initially calculated independently for each B-AR, as opposed to summed across all ratios.

Statistical Analysis

All statistical analyses were performed with IBM SPSS Statistics 23.0 (IBM, Armonk, NY). Descriptive statistics were first calculated for all variables and the means for all variables were plotted by B-AR to examine the response of each to different ratios. Ball-to-aperture ratios were eliminated based on visual inspection of plots, and variables were recalculated, summed across all remaining ratios, for analysis. Intraclass correlation coefficients [ICC (3,1)] were calculated for all variables for one and two cycles of testing to determine the intersession reliability. Coefficients of variation were calculated using the typical error of

the measure (CV_{TE}), as described by Hopkins.^{13,14} Log transformations were applied to MT before calculating CV_{TE} , as the variable showed significant departures from normality.¹³ Level for statistical significance was set a priori at: $\alpha = 0.05$.

RESULTS

Visual inspection of plots for the effect of B-AR on PACT variables (Fig. 2) showed a typical, quadratic pattern with RT, MT, IT, and RST increasing as B-AR approached 1.0, and ACC decreasing. It also appeared that successive B-ARs furthest from 1.0 (0.2–0.4 and 1.6–1.8) elicited fairly similar mean responses for most variables (mean difference = $1.3 \pm 1.2\%$).

Subsequent sets of successive ratios, starting with 0.4–0.6 and 1.4–1.6 (mean difference = $2.1 \pm 2.4\%$), started to show larger differences. Therefore, only the furthest ratios of 0.2 and 1.8 were eliminated. Moving up one B-AR closer to 1.0 (0.4–0.6 and 1.4–1.6), it did not appear that these successive ratios elicited similar mean responses for most variables, with timing variables beginning to increase slightly and ACC beginning to decrease.

All results for ICCs and CV_{TE} are presented in Table I. Overall, all variables showed good to excellent intersession reliability with both 2 (ICC = 0.808–0.985) and 1 (ICC = 0.709–0.937) cycles of testing. Further, all ICCs were statistically significant ($P < 0.05$). For within-subject stability, mean CV_{TE} ranged from 2.18–14.67% with 2 cycles of testing, and from 3.39–19.26% for 1 cycle of testing.

EXPERIMENT 2

Each cycle of the PACT consists of 16 presentations of each B-AR (Sec. 2.1), in a block-randomized order, for a total of 128 trials. This large number of trials was selected for the initial version of the PACT to ensure that enough trials were completed to capture perceptual-motor decision making during preliminary work. Establishing the minimum number of trials necessary to capture the intended behavioral response from subjects is imperative, given that performing more trials than necessary can alter observed response behaviors due to boredom and mental fatigue. Therefore, the purpose of Experiment 2 was to test the effects of eliminating trials on the intersession reliability and within-subject stability of the PACT. Not only would this serve to limit the required testing further, but also to establish the threshold number of trials necessary to obtain adequate reliability and stability across all variables.




Fig. 2. Means for all PACT variables by ball-to-aperture ratio.

A secondary purpose was to establish these same parameters, for all variables, within each individual B-AR. As discussed previously (Sec. 1.0), accuracy and reaction times of affordance judgments are known to be affected by the proximity of the stimulus to the relevant action boundary.^{18,21,22} Further, the results of Experiment 1 confirmed this effect for the PACT, with B-ARs approaching 1.0 eliciting slower, less accurate responses. Subsequently, researchers utilizing the PACT in future studies may be interested in how individuals perform on the PACT in response to specific B-ARs. It may be that environmental, physiological or cognitive stressors will start to elicit changes in performance closer to the action boundary before overall performance begins to deteriorate or will only elicit changes closer to the action boundary. Therefore, we sought to also provide information on the threshold number of trials necessary to obtain adequate reliability and stability for PACT variables for each individual B-AR.

METHODS

Subjects and Procedures

Data collected from the same longitudinal study was utilized for Experiment 2. All PACT variables (RT, MT, IT, RST, ACC) were calculated as described above, averaged across the 4th and 5th cycles of each session, and with the 0.2 and 1.8 B-ARs removed.

Data reduction was iterative, beginning with 32 trials (2 cycles of 16 trials each) of each B-AR, from which 4 trials of each B-AR were eliminated and PACT variables were re-averaged across all remaining trials (i.e., 32, 28, 24, ..., 4 trials). In total, 24 trials were eliminated at each step. This process was utilized in recalculating all variables averaged across all B-AR, as well as within each B-AR. During the PACT, each B-AR is presented in a block-randomized order, meaning each ratio is presented once before any are presented twice. This allowed for the elimination of an equal number of trials for each B-AR,

Table I. Intraclass Correlation Coefficients and Coefficients of Variation for 2 and 1 Cycles of Testing and With 0.2 and 1.8 Ball-to-Aperture Ratios Removed.

	ICC*	95% CI	MEAN CV _{TE} (%)	95% CI
Reaction Time				
2 Cycles	0.828	0.629–0.933	12.83	10.42–16.76
1 Cycle	0.824	0.623–0.932	11.93	9.69–15.59
Movement Time				
2 Cycles	0.985	0.968–0.994	14.67	11.76–19.59
1 Cycle	0.940	0.873–0.977	19.26	15.38–25.87
Initiation Time				
2 Cycles	0.934	0.844–0.975	10.71	8.70–14.00
1 Cycle	0.899	0.768–0.962	12.05	9.79–15.74
Response Time				
2 Cycles	0.918	0.825–0.968	6.61	5.37–8.63
1 Cycle	0.870	0.722–0.949	8.03	6.52–10.49
Accuracy				
2 Cycles	0.808	0.595–0.924	2.18	1.77–2.84
1 Cycle	0.709	0.394–0.885	3.39	2.75–4.43

* ICC = intraclass correlation coefficient; 95% CI = 95% confidence interval; CV_{TE} = coefficient of variation using typical error.

while still maintaining the sequential order in which trials were presented. In simpler terms, when the first 4 trials were eliminated for each B-AR, the remaining 28 trials were trials 1–28, in chronological order.

Statistical Analysis

Descriptive statistics were first calculated for all variables. Intraclass correlation coefficients [ICC (3,1)] were calculated for all variables for each iteration of eliminating trials to determine the intersession reliability. Coefficients of variation were calculated using the typical error of the measure, and a log transformation was applied to all MT coefficients.¹⁴ Finally, means for all variables were plotted by B-AR, based on the results of these analyses, to re-examine the continuum of responses about the action boundary.

RESULTS

Results of ICCs and CV_{TE} with variables averaged across ratios are reported in **Table II**. For RT, MT, IT, and RST, ICCs showed good to excellent reliability until the number of remaining trials for each B-AR reached 20 (ICC = 0.817–0.973). For ACC, ICCs showed good reliability until the number of remaining trials reached 24 (ICC = 0.811). In regards to within-subject stability, CV_{TE} remained relatively stable for RT through all trials (Mean CV_{TE} = 10.89–12.83%) and for ACC through 4 trials (mean CV_{TE} = 2.12–3.82%). Initiation time (mean CV_{TE} = 10.08–12.05%) and RST (mean CV_{TE} = 6.57–8.03%) both remained relatively stable through 16 trials. Finally, MT showed the highest CV_{TE} and remained stable through 20 trials (mean CV_{TE} = 14.67–15.61%).

Results of ICCs and CV_{TE} with variables averaged for each individual B-AR are presented in **Tables III** and **IV**. To limit the size of tables, results are only reported for averages of 20 trials or greater, given the results for variables averaged across ratios.

Table II. Intraclass Correlation Coefficients and Coefficients of Variation: Demonstrating the Effects of Incrementally Removing Four Trials of Each Ratio.

	ICC*	95% CI	MEAN CV _{TE} (%)	95% CI
Reaction Time				
32 Trials	0.828	0.629–0.933	12.83	10.42–16.76
28 Trials	0.841	0.658–0.938	11.08	9.00–14.47
24 Trials	0.822	0.620–0.931	10.89	8.85–14.23
20 Trials	0.817	0.608–0.929	11.98	9.73–15.65
16 Trials	0.824	0.623–0.932	11.93	9.69–15.59
12 Trials	0.832	0.640–0.935	11.55	9.38–15.09
8 Trials	0.828	0.629–0.933	12.02	9.77–15.71
4 Trials	0.846	0.669–0.940	11.55	9.38–15.08
Movement Time				
32 Trials	0.985	0.968–0.994	14.67	11.76–19.59
28 Trials	0.972	0.941–0.989	15.45	12.38–20.65
24 Trials	0.974	0.945–0.990	15.61	12.50–20.86
20 Trials	0.973	0.943–0.990	15.40	12.34–20.57
16 Trials	0.940	0.873–0.977	19.26	15.38–25.87
12 Trials	0.919	0.828–0.969	23.46	18.67–31.70
8 Trials	0.897	0.781–0.960	27.88	22.11–37.88
4 Trials	0.868	0.719–0.948	33.74	26.64–46.20
Initiation Time				
32 Trials	0.934	0.844–0.975	10.71	8.70–14.00
28 Trials	0.932	0.839–0.975	11.29	9.17–14.75
24 Trials	0.933	0.840–0.975	10.96	8.91–14.32
20 Trials	0.935	0.841–0.976	10.08	8.19–13.17
16 Trials	0.899	0.768–0.962	12.05	9.79–15.74
12 Trials	0.874	0.718–0.951	14.29	11.61–18.67
8 Trials	0.853	0.673–0.943	16.06	13.05–20.98
4 Trials	0.841	0.651–0.938	16.91	13.74–22.09
Response Time				
32 Trials	0.918	0.825–0.968	6.61	5.37–8.63
28 Trials	0.918	0.819–0.968	6.63	5.38–8.66
24 Trials	0.920	0.824–0.969	6.57	5.34–8.58
20 Trials	0.918	0.818–0.968	7.26	5.90–9.49
16 Trials	0.870	0.722–0.949	8.03	6.52–10.49
12 Trials	0.833	0.647–0.934	9.08	7.38–11.87
8 Trials	0.815	0.610–0.927	9.31	7.56–12.17
4 Trials	0.737	0.454–0.896	10.85	8.81–14.18
Accuracy				
32 Trials	0.808	0.595–0.924	2.18	1.77–2.84
28 Trials	0.849	0.682–0.941	2.12	1.73–2.77
24 Trials	0.811	0.595–0.926	2.57	2.09–3.36
20 Trials	0.752	0.462–0.904	2.90	2.35–3.78
16 Trials	0.709	0.394–0.885	3.39	2.75–4.43
12 Trials	0.685	0.344–0.875	3.71	3.02–4.85
8 Trials	0.398	-0.198–0.755	3.82	3.10–4.99
4 Trials	0.336	-0.252–0.721	6.33	5.14–8.27

* ICC = intraclass correlation coefficient; 95% CI = 95% confidence interval; CV_{TE} = coefficient of variation using typical error.

Across all variables except ACC, all B-ARs showed similar results for intersession reliability with 24 trials of testing, compared to averaging variables across ratios, with ICCs ≥ 0.776 . For ACC, it appears that reliability is reduced when looking at individual B-ARs, with only ratios of 1.4 (ICC = 0.726–0.797) and 1.6 (ICC = 0.668–0.720) showing similar results compared to averaging ACC across ratios. Coefficients of variation were similar for all B-ARs for IT, RST, and ACC compared to averaging variables across ratios. However, CV_{TE} for RT and MT showed marked increases for most ratios. Finally, plots depicting means for all variables by B-AR, after removal of ratios and trials (24 trials remaining), are shown in **Fig. 3**.

Table III. Intraclass Correlation Coefficients Within Each Ball-to-Aperture Ratio.*

	0.4	0.6	0.8	1.2	1.4	1.6
Reaction Time						
32 Trials	0.872	0.824	0.793	0.693	0.815	0.830
28 Trials	0.879	0.827	0.807	0.713	0.833	0.837
24 Trials	0.861	0.807	0.776	0.685	0.814	0.831
20 Trials	0.831	0.779	0.671	0.588	0.752	0.813
Movement Time						
32 Trials	0.971	0.968	0.965	0.973	0.971	0.963
28 Trials	0.969	0.965	0.961	0.970	0.067	0.961
24 Trials	0.970	0.962	0.956	0.972	0.969	0.964
20 Trials	0.965	0.946	0.954	0.968	0.970	0.953
Initiation Time						
32 Trials	0.916	0.928	0.902	0.920	0.938	0.945
28 Trials	0.917	0.920	0.895	0.918	0.938	0.941
24 Trials	0.916	0.921	0.888	0.922	0.933	0.943
20 Trials	0.904	0.904	0.879	0.935	0.932	0.948
Response Time						
32 Trials	0.930	0.905	0.856	0.880	0.910	0.912
28 Trials	0.930	0.901	0.861	0.859	0.894	0.914
24 Trials	0.931	0.892	0.833	0.850	0.896	0.921
20 Trials	0.910	0.855	0.844	0.850	0.865	0.889
Accuracy						
32 Trials	0.590	0.621	0.657	0.547	0.811	0.666
28 Trials	0.565	0.638	0.681	0.615	0.822	0.711
24 Trials	0.423	0.607	0.696	0.578	0.797	0.668
20 Trials	0.435	0.494	0.544	0.357	0.726	0.720

* All data are results of intraclass correlation coefficients [ICC (3,1)].

DISCUSSION

The PACT is a novel and ecologically valid assessment of perceptual-motor accuracy and reaction times.^{7,9,21} The purpose of the current study was to explore several methods to reduce PACT testing time while maintaining the intersession reliability and within-subject stability previously established for this assessment. In this effort, two experiments were undertaken. For Experiment 1, the relationship of reaction times and judgment accuracy on the PACT with B-ARs was first assessed, and based on these relationships, the ratios of 0.2 and 1.8 were eliminated. Variables were recalculated, and the effect of removing these two ratios was assessed. For Experiment 2, this same process was repeated after eliminating blocks of trials and reliability and within-subject stability coefficients were also calculated for individual B-ARs.

The results of Experiment 1 demonstrated that the ratios furthest from the action boundary could be eliminated from PACT testing without producing a large effect on either intersession reliability or within-subject stability. In the previous study by Connaboy *et al.*,⁷ ICCs for one cycle of testing and with all B-ARs included ranged from 0.707–0.943. In comparison, IT showed the largest negative effect on reliability with the elimination of ratios, decreasing from 0.906 to 0.899. Likewise, ICCs for two cycles of testing and with all B-ARs were reported to range from 0.820–0.992. Again, IT showed the largest negative effect, decreasing from 0.992 to 0.934. More importantly, all variables retained good to excellent reliability with two cycles of testing, according to a range of commonly cited criteria for the

Table IV. Mean Coefficients of Variation Within Each Ball-to-Aperture Ratio.

	0.4 (%)	0.6 (%)	0.8 (%)	1.2 (%)	1.4 (%)	1.6 (%)
Reaction Time						
32 Trials	12.74	14.36	13.98	14.20	13.29	12.67
28 Trials	12.74	13.80	14.22	14.52	13.56	13.64
24 Trials	12.76	14.40	15.60	15.53	13.84	13.98
20 Trials	15.98	15.49	17.39	17.42	14.20	14.27
Movement Time						
32 Trials	18.67	16.35	20.55	20.29	18.16	23.62
28 Trials	19.48	16.36	21.95	20.30	19.05	23.78
24 Trials	22.53	18.82	22.20	19.95	21.08	24.18
20 Trials	25.30	21.28	22.36	20.50	21.05	28.54
Initiation Time						
32 Trials	11.93	11.50	13.05	12.64	11.02	10.46
28 Trials	12.08	12.31	13.62	12.52	10.86	10.87
24 Trials	12.05	11.93	14.32	11.86	11.62	10.02
20 Trials	11.94	12.69	14.47	10.92	11.35	9.59
Response Time						
32 Trials	6.81	7.19	7.76	7.57	7.26	8.41
28 Trials	6.96	7.21	7.73	8.09	8.21	8.73
24 Trials	7.40	7.84	8.11	8.45	8.32	9.08
20 Trials	8.95	9.21	8.66	8.56	8.52	11.25
Accuracy						
32 Trials	2.94	3.11	5.46	4.37	3.60	3.36
28 Trials	3.22	3.40	5.36	4.51	3.45	3.80
24 Trials	3.80	4.46	5.74	5.54	3.87	4.48
20 Trials	3.10	5.19	6.84	6.43	4.61	5.26

All data are results of mean coefficients of variation calculated using the typical error of the mean.

interpretation of reliability coefficients.^{2,12,19} Further, most variables showed good reliability with only one cycle of testing, however ACC may require 2 (ICC = 0.709 with 1 cycle) depending on the criteria used.^{2,12,19}

Coefficients of variation with all B-ARs were previously reported to range from 2.05–16.03% with two cycles of testing.⁷ Initiation time showed the largest increase in the CV_{TE} with elimination of ratios, moving from 4.07 to 10.71%. For one cycle of testing, previously reported CV_{TE} ranged from 3.28–19.28%, with IT also showing the largest increase with elimination of ratios, moving from 11.88 to 12.05%. With only IT showing a marked increase in CV_{TE}, and only for two cycles of testing, it appears that within-subject stability remained stable with the removal of ratios. In fact, for several variables (MT and RT) the CV_{TE} were decreased and the ICCs increased, compared to those previously reported, indicating improved stability and reliability.⁷

Based on the results of Experiment 1, Experiment 2 moved forward with only B-ARs of 0.4–1.6, while retaining two cycles of testing. The primary results of Experiment 2 demonstrated that 8–12 trials of each B-AR could be eliminated from PACT testing while still maintaining the intersession reliability and within-subject stability of all variables. All variables maintained good to excellent intersession reliability with the elimination of 12 trials (ICCs = 0.817–0.973) except for ACC (ICC = 0.752). However, ACC maintained good reliability through the elimination of eight trials (ICC = 0.811). Within-subject stability results mirrored those for reliability, with CV_{TE} remaining relatively unchanged through the elimination of 12 trials for all




Fig. 3. Means for all PACT variables by ball-to-aperture ratio after removal of 2 ratios and 12 trials from each remaining ratio.

variables ($CV_{TE} = 2.90\text{--}15.40\%$). With the elimination of further trials, CV_{TE} showed a large increase for several variables, including MT (15.40–19.26%) and IT (10.08–12.05%).

Finally, PACT variables still show similar or superior inter-session reliability and stability, after the elimination of 2 B-ARs and 12 trials for each ratio, compared to other cognitive measures of a similar nature (choice/procedural reaction time measures). Previous literature has reported ICCs ranging from 0.26–0.69 for different cognitive performance measures.^{3,6,20} In contrast, the PACT variable with the lowest reliability coefficient still showed an ICC of 0.752. Comparisons for within-subject stability are more difficult, as the majority of previous studies do not report coefficients of variation. However, two studies that have included the standard difference of the error for choice/procedural reaction time tasks, a similar measure

of within-subject stability, reported values of 12.77% and 19.38%.^{6,20} In comparison, CV_{TE} for all PACT variables (2.90–15.40%) were within this range or lower.

A secondary purpose of Experiment 2 was to examine ICCs and CV_{TE} for PACT variables, averaged for each independent B-AR. This information may be useful for investigators interested in the effects of a given perturbation or intervention on responses to specific B-ARs, or on the continuum of responses about the action boundary.^{18,21,22} Overall, the results of these tests showed that reliability was decreased for ACC, and within-subject variability was increased for RT and MT, compared to averaging variables across all B-ARs. While this may be of concern, it is worth noting that ICCs and CV_{TE} were comparable to similar cognitive measures.^{3,6,20} The addition of further trials (up to 32) resulted in improvements for several

variables, within several B-ARs, however most remained unchanged. Finally, plots for all variables by B-AR with 24 trials for each ratio (Fig. 3) showed similar patterns as those before trials were removed (Fig. 2). Initiation time appeared to have a slightly flatter curve, however this variable also showed the flat-test pattern about the action boundary before the elimination of trials.

In conclusion, the combined results of Experiment 1 and 2 demonstrate that all PACT variables are reliable and stable when B-ARs are limited to 0.4–1.6, and only 24 trials of each B-AR are retained. This represents an almost 45% reduction in total testing time, from 256 trials to 144. Further, over the 3-cycle familiarization period, this would result in a 50% reduction in testing time, from 384 trials to 192. In more practical terms, this means that the shortened PACT would require a familiarization period of approximately 7.5 min, and a testing period of 5.5 min; down from 15 and 10 min, respectively. These changes, therefore, would hold great significance in reducing the time-burden of the PACT. When working with pilots, astronauts, or other military personnel, subjects or patients are often under stringent time constraints. Therefore, these reductions in testing burden are highly significant for researchers or clinicians using the PACT to evaluate behavioral risk in these populations.

A limitation of the current study is the extrapolation of previously collected data to these conclusions. While the time sequence for performance of trials was maintained when eliminating ratios and trials, the cycles eliminated for familiarization were performed with all trials and ratios. Therefore, we cannot say whether these results would hold if all cycles were performed with the limited ratios and trials outlined above. However, the purpose of this study was to investigate these changes to PACT testing and outline the best course for its future development. In this sense, we believe that these results still hold great significance in providing evidence for the efficacy of a shortened PACT, even if future work is needed to confirm our findings.

ACKNOWLEDGMENTS

The opinions or assertions contained herein are the views of the authors and are not to be constructed as official or reflecting the views of NASA. Further, the funding body did not have a role in the design, analysis, or interpretation of the current study.

Financial Disclosure Statement: This study was funded by the National Aeronautics and Space Administration; grant number NNX15AC13G. The authors have no conflicts to declare.

Authors and affiliations: Caleb D. Johnson, Ph.D., Alice D. LaGoy, M.S., B.Sc., Shawn R. Eagle, Ph.D., Anne A. Beethe, M.A., B.Sc., and Christopher Connaboy, Ph.D., Neuromuscular Research Laboratory, Department of Sports Medicine and Nutrition, University of Pittsburgh, Pittsburgh, PA; Gert-Jan Pepping, Ph.D., School of Exercise Science, Australian Catholic University, Brisbane Campus, Banyo, Australia; Joanne L. Bower, Ph.D., Department of Psychology, De Montfort University, The Newarke, Leicester, UK; Candice A. Alfano, Ph.D., Department of Psychology, University of Houston, Houston, TX; and Richard J. Simpson, Ph.D., Department of Nutritional Sciences, The University of Arizona, Tucson, AZ.

REFERENCES

1. Aherne BB, Zhang C, Chen WS, Newman DG. Pilot decision making in weather-related night fatal helicopter emergency medical service accidents. *Aerosp Med Hum Perform.* 2018; 89(9):830–836.
2. Anastasi A. *Psychological testing.* New York (NY): Collier Macmillan; 1988.
3. Ayala F, De Ste Croix M, Sainz de Baranda P, Santonja F. Inter-session reliability and sex-related differences in hamstrings total reaction time, pre-motor time and motor time during eccentric isokinetic contractions in recreational athlete. *J Electromyogr Kinesiol.* 2014; 24(2):200–206.
4. Bouak F, Vartanian O, Hofer K, Cheung B. Acute mild hypoxic hypoxia effects on cognitive and simulated aircraft pilot performance. *Aerosp Med Hum Perform.* 2018; 89(6):526–535.
5. Chmura J, Krysztofiak H, Ziemba AW, Nazar K, Kaciuba-Uscilko H. Psychomotor performance during prolonged exercise above and below the blood lactate threshold. *Eur J Appl Physiol Occup Physiol.* 1998; 77(1–2):77–80.
6. Cole WR, Arriex JP, Schwab K, Ivins BJ, Qashu FM, Lewis SC. Test-retest reliability of four computerized neurocognitive assessment tools in an active duty military population. *Arch Clin Neuropsychol.* 2013; 28(7):732–742.
7. Connaboy C, Johnson CD, LaGoy AD, Pepping G-J, Simpson RJ, et al. Intersession reliability and within-session stability of a novel perception-action coupling task. *Aerosp Med Hum Perform.* 2019; 90(2):77–83.
8. Daviaux Y, Mignardot J-B, Cornu C, Deschamps T. Effects of total sleep deprivation on the perception of action capabilities. *Exp Brain Res.* 2014; 232(7):2243–2253.
9. Eagle SR, Nindl BC, Johnson CD, Kontos AP, Connaboy C. Does concussion affect perception-action coupling behavior?: Action boundary perception as a biomarker for concussion. *Clin J Sport Med.* 2019. [Epub ahead of print.]
10. Fajen BR, Riley MA, Turvey MT. Information, affordances, and the control of action in sport. *Int J Sport Psychol.* 2009; 40(1):79.
11. Gibson James J. The theory of affordances. In: Shaw R, Bransford J, editors. *Perceiving, Acting, and Knowing.* Hoboken (NJ): John Wiley & Sons; 1977.
12. Heaton RK, Temkin N, Dikmen S, Avitable N, Taylor MJ, et al. Detecting change: A comparison of three neuropsychological methods, using normal and clinical samples. *Arch Clin Neuropsychol.* 2001; 16(1):75–91.
13. Hopkins WG. Measures of reliability in sports medicine and science. *Sports Med.* 2000; 30(1):1–15.
14. Hopkins WG. Spreadsheets for analysis of validity and reliability. A new view of statistics. *Sportscience* 2015; 19:36–42.
15. Kopp B, Rist F, Mattler U. N200 in the flanker task as a neurobehavioral tool for investigating executive control. *Psychophysiology.* 1996; 33(3):282–294.
16. Logan GD, Cowan WB, Davis KA. On the ability to inhibit simple and choice reaction time responses: a model and a method. *J Exp Psychol Hum Percept Perform.* 1984; 10(2):276–291.
17. Palmer CJ, Bigelow C, Van Emmerik RE. Defining soldier equipment trade space: load effects on combat marksmanship and perception-action coupling. *Ergonomics.* 2013; 56(11):1708–1721.
18. Pepping G-J, Li F-X. Effects of response task on reaction time and the detection of affordances. *Motor Control.* 2005; 9(2):129–143.
19. Portney L, Watkins M. *Foundations of clinical research: application to practice.* Stamford (CT): Appleton & Lange; 1993.
20. Register-Mihalik JK, Guskiewicz KM, Mihalik JP, Schmidt JD, Kerr ZY, McCrea MA. Reliable change, sensitivity, and specificity of a multidimensional concussion assessment battery: Implications for caution in clinical practice. *J Head Trauma Rehabil.* 2013; 28(4):274–283.

21. Smith J, Pepping G-J. Effects of affordance perception on the initiation and actualization of action. *Ecol Psychol.* 2010; 22(2):119–149.
22. Stins JF, Michaels CF. Stimulus-response compatibility for absolute and relative spatial correspondence in reaching and in button pressing. *Q J Exp Psychol A.* 2000; 53(2):569–589.
23. Vrijkotte S, Roelands B, Meeusen R, Pattyn N. Sustained military operations and cognitive performance. *Aerosp Med Hum Perform.* 2016; 87(8):718–727.
24. Withagen R, De Poel HJ, Araújo D, Pepping G-J. Affordances can invite behavior: Reconsidering the relationship between affordances and agency. *New Ideas Psychol.* 2012; 30(2):250–258.