

Temporary Incapacitation Rates and Permanent Loss of Medical License in Commercial Airline Pilots

Erik Hohmann; Reino Pieterse

INTRODUCTION: The purpose of this study was to report the temporary loss of medical license and pilot incapacitations in the United Arab Emirates from 2018–2021.

METHOD: The General Civil Aviation Authority database was searched for all reported temporary suspensions of license between 2018–2021 and the ICD-10 codes were extracted.

RESULTS: A total of 1233 incapacitations was reported with a mean license suspension of 148.4 ± 276.8 d. The mean days of suspension for the various medical specialties were 115.2 ± 188.4 for musculoskeletal conditions ($N = 392$), 189.3 ± 324.8 for medicine ($N = 335$), 101.6 ± 231.4 for surgery, 109.4 ± 223.5 for urology ($N = 93$), 90.3 ± 128.7 for ophthalmology ($N = 68$), 385.6 ± 594.3 for psychiatry ($N = 61$), 150.4 ± 285.9 for ENT ($N = 59$), 419.4 ± 382.6 for obstetrics and gynecology ($N = 30$), and 44.9 ± 39 for dermatology ($N = 21$). Permanent suspensions were as follows: total $N = 100$ (8.1%), musculoskeletal $N = 13$ (3.3%), medicine $N = 37$ (11%), surgery $N = 10$ (5.7%), urology $N = 10$ (10.7%), ophthalmology $N = 2$ (2.9%), psychiatry $N = 20$ (32.8%), ENT $N = 1$ (1.7%), obstetrics and gynecology $N = 4$ (13.1%), and dermatology $N = 3$ (14.3%).

DISCUSSION: Musculoskeletal conditions are the most common reason for temporary loss of medical license followed by medical and surgical conditions. The least common reason was dermatological conditions. The longest period of incapacitation was associated with psychiatric conditions followed by medical and ENT conditions. The annual calculated temporary incapacitation rate was 2.8% and the permanent suspension rate was 0.25%.

KEYWORDS: return to work, pilots, temporary medical license suspension, occupational health.

Hohmann E, Pieterse R. *Temporary incapacitation rates and permanent loss of medical license in commercial airline pilots. Aerosp Med Hum Perform.* 2023;94(3):113–121.

Work and working conditions have a profound effect on an individual's health.² Perceived job insecurity, physical exposures to dangerous conditions, but also rotating shift work can result in mental and physical conditions and deteriorating health.² For example, shift work and long work hours have the potential to increase the risk of injury and obesity and result in a wide range of chronic diseases.⁵ In long-haul truck drivers the risk for cardiovascular and metabolic disease was 3.7 times, respectively 4.31 higher when compared to the National Health and Nutrition Examination Survey sample of the general U.S. population.^{14,22}

Airline pilots work in similar conditions which involve shift-work, circadian rhythm disturbances with exposure to mild hypoxia, reduced atmospheric pressure, low humidity, noise vibration, and cosmic radiation.³ Butler and Nicholas performed a survey of more than 10,000 active and retired pilots in

the United States and Canada and reported that there is an increased risk of melanoma, motor neuron disease, and cataracts.³ However, the authors could not demonstrate an increased risk for other diseases such as heart disease, diabetes, and high blood pressure when comparing their data to the 1995 National Health Interview Survey.³ Evans and Radcliffe have reported annual incapacitation rates in commercial airline pilots and

From the Burjeel Hospital for Advanced Surgery and the Emirates Group Rehabilitation Unit, Dubai, United Arab Emirates.

This manuscript was received for review in February 2022. It was accepted for publication in January 2023.

Address correspondence to: Erik Hohmann, School of Medicine, Faculty of Health Sciences, University of Pretoria, Cnr Bophelo and Dr. Savage Road, Gezina, Pretoria 0001, South Africa; ehohmann@hotmail.com.

Reprint and copyright © by the Aerospace Medical Association, Alexandria, VA.

DOI: <https://doi.org/10.3357/AMHP.6071.2023>

have shown that accidents, cardiovascular, and musculoskeletal conditions were the most common reasons for temporary unfitness.⁸ Parker and Snyder showed that 25% of airline pilot morbidity and long-term disability occurred because of cardiovascular reasons with no correlation to age followed by 11% for orthopedic and musculoskeletal conditions.⁸ Long-term disability rates in a cohort of 2271 Canadian pilots were recorded between 1981 and 1990.^{1,22} The main reason for a long duration of leave of absence were mental disorders (16%), circulatory (14%) and digestive (12%) system diseases, and musculoskeletal conditions (11%).^{1,22} Currently there is insufficient updated information on medical conditions resulting in temporary loss of medical license and incapacitation. The purpose of this study was to therefore report the temporary loss of medical license and pilot incapacitations in the United Arab Emirates from 2018–2021.

METHODS

This study used data from the database of the General Civil Aviation Authority (GCAA) and the following data was extracted by an aeromedical officer of the GCAA: date of license suspension and reinstatement, age, gender, and ICD-10 codes. The data was then forwarded to the first author. The authors had no access to any other information and this process effectively de-identified personal data.⁴ In general, registry data studies are exempt from IRB approval and our study design is in compliance with the World Medical Association declaration of Helsinki ethical principles for medical research amended by the 64th WMA General Assembly in 2013.¹⁹ Admittedly, the available research data could be de-identified, but the likelihood of unauthorized access to the GCAA database by a non-GCAA employee is not possible.

The GCAA is the federal regulatory body of the United Arab Emirates, overseeing all aviation-related activities in the United Arab Emirates. The GCAA CAR MED regulations outline the medical provisions for licensing pilots and cabin crew.¹¹ If pilots were determined to be unfit, the responsible aviation medical examiner will upload a request for suspension of the medical license onto the GCAA website.¹⁶ Once the report is uploaded onto the GCAA website, the license will be suspended with immediate effect. For medical conditions, a medical report will also be submitted to the GCAA which includes all applicable ICD-10 codes. Similarly, for reinstatement of an aviation license, the responsible aviation medical physician will upload a request for reinstatement. This request again includes all applicable ICD-10 codes, all applicable medical reports, and detailed operation notes from the treating specialist.¹⁶

The following inclusion criteria were applied: license suspension for all pilots registered with the GCAA between January 2018 to October 2021, age between 18–65 yr, and availability of ICD-10 codes for all license suspensions. Pilots were excluded if they voluntarily suspended their license. If there was no date of reinstatement, pilots were considered to

have lost their license permanently. Medical and surgical pathology was categorized according to the ICD-1-PCS system (ICD10Data.com) and the following categories were used: dermatology, ear-nose-throat (ENT), obstetrics and gynecology, musculoskeletal, ophthalmology, psychiatry, surgical, and urology.

Statistical Analysis

Descriptive analysis was applied to demographic data. The mean, standard deviations, ranges, and 95% confidence intervals for the time of pilot incapacitation were calculated for the total number of cases. The data was then subdivided into nine subspecialties: musculoskeletal, medicine, surgical, urology, ophthalmology, psychiatry, ENT, obstetrics and gynecology, and dermatology.

For the individual subspecialties, similar pathology and ICD-10 codes were grouped and means and standard deviations were calculated. Pilots who did not return to work and had permanent license suspensions were excluded when calculating mean and standard deviation with subgroup pooling. All analyses were conducted using STATA SE for Windows (version 12.0; StataCorp, College Station, TX, USA).

RESULTS

Between January 2018 to October 2021, a total of 1233 pilots was reported to the GCAA for medical incapacity and their license was temporarily suspended. The mean age was 46.4 ± 10.4 yr. There were 1173 male pilots with a mean age of 46.9 ± 10.3 yr and 60 female pilots with a mean age of 37.1 ± 7.7 yr. The mean days of temporary suspension were 148.4 ± 276.8 d, ranging from 1–1257 d with a 95% confidence interval between 63–98.6 d. The largest number of pilot incapacitations was reported for musculoskeletal conditions and the lowest number was reported for dermatological conditions. The longest time of incapacitation was observed for pilots with psychiatric conditions and the shortest time was reported for pilots with dermatological conditions. Of the pilots, 100 did not return to work. **Table I** summarizes the period of incapacitation for the study period.

Musculoskeletal

There was a total of 392 pilots with medical conditions (**Table II**). The mean age was 46.2 ± 9.3 yr. There were 383 men with a mean age of 46.2 ± 9.4 and 9 women with a mean age of 47.6 ± 7 yr. The most common conditions were meniscus injuries ($N = 39$ with a mean suspension of 56.8 d), lumbar spine disc prolapse ($N = 23$ with a mean suspension of 184.5 d), and shoulder impingement ($N = 19$ with a mean suspension of 121.1 d). However, the most common reason for suspension was orthopedic trauma, with a total of 131 cases. The mean time of suspension was 115.2 ± 188.4 d. There were 13 pilots who did not return to work. The reasons were related to fractures in six cases, spinal conditions in three cases, and osteoarthritis and other degenerative conditions in four cases.

Table I. Period of Temporary Incapacitation for Pilots.

	N	MEAN DAYS	SD	RANGE	95% CI	DID NOT RETURN
Musculoskeletal	392	115.2	188.4	2–1446	96.3–134.2	13
Medicine	335	189.3	324.8	58–2170	151.7–226.9	37
Surgical	174	101.6	231.4	1–2025	64.9–138.4	10
Urology	93	109.4	223.5	2–1268	27.8–131.5	10
Ophthalmology	68	90.3	128.7	2–719	58.6–121.9	2
Psychiatry	61	385.6	594.3	6–2757	195.5–575.6	20
ENT	59	150.4	285.9	1–1727	75.3–225.6	1
Obstetrics/Gynecology	30	419.4	382.6	5–1390	268–572.7	4
Dermatology	21	44.9	39.6	9–162	25.2–64.6	3
Total	1233	148.4	276.8	1–2757	63.0–98.6	100

Medicine

There was a total of 335 pilots with medical conditions (Table III). The mean age was 49.5 ± 9.7 yr. There were 333 men with a mean age of 49.6 ± 9.7 yr and 2 women with a mean age of 39.5 ± 4.9 yr. The most common conditions were hypertension ($N = 60$ with a mean suspension of 101.6 d), sleep apnea ($N = 28$ with a mean suspension of 173 d), angina ($N = 27$ with a mean suspension of 217.5 d), diabetes mellitus ($N = 26$ with a mean suspension of 116.9 d), and arrhythmias ($N = 26$ with a mean suspension of 170.6 d). The mean time of suspension was 189.3 ± 324.8 d. A total of 37 pilots did not return to work. The reasons were related to cardiac conditions in 10 cases, pulmonary conditions in 4 cases, gastroenterological conditions in 2 cases, neurological conditions in 9 cases, infectious conditions in 1 case (malaria), malignancies in 7 cases, and endocrinological conditions in 2 cases.

Surgery

There was a total of 174 pilots with surgical conditions (Table IV). The mean age was 45.3 ± 9.8 yr. There were 171 men with a mean age of 45.5 ± 9.7 yr and 3 women with a mean age of 32.7 ± 5.8 yr. The most common conditions were inguinal hernia ($N = 53$ with a mean suspension of 48.4 d), appendicitis ($N = 24$ with a mean suspension of 35.8 d), and cholecystitis ($N = 17$ with a mean suspension of 120.1 d). The mean time of suspension was 101.6 ± 231.4 d. Of the 174, 10 pilots did not return to work. Two pilots did not return following appendicitis surgery and one pilot each did not return for the following conditions: aortic aneurysm, concussion, bariatric surgery, following surgery for cholecystitis, following surgery for inguinal hernia, pilonidal sinus, stricture of artery, and following thyroid surgery.

Urology

There was a total of 93 pilots with urological conditions (Table V). The mean age was 46.8 ± 10.5 yr. There were 90 men with a mean age of 47.1 ± 10.6 yr and 3 women with a mean age of 37.7 ± 3.1 yr. The most common conditions were kidney stones ($N = 35$ with a mean suspension of 96.6 d) and ureter stones ($N = 26$ with a mean suspension of 99.5 d). The mean time of suspension was 109.4 ± 223.5 d. A total of 10 pilots did not return to work. Five pilots were diagnosed with prostate cancer, two pilots with bladder cancer, one pilot with kidney

cancer, one pilot with testicular cancer, and one pilot did not return following ureter stones.

Ophthalmology

There was a total of 68 pilots with ophthalmological conditions (Table VI). The mean age was 44.8 ± 13.2 yr. There were 67 men with a mean age of 45.1 ± 13.1 and 1 woman who was 23 yr old. The most common conditions were cataract surgery ($N = 14$ with a mean suspension of 136 d) and retina disorders ($N = 14$ with a mean suspension of 106.6 d). The mean time of suspension was 90.3 ± 128.7 d. Two pilots did not return to work; one following cataract surgery and one who was diagnosed with glaucoma.

Psychiatry

There was a total of 61 pilots with psychiatric conditions (Table VII). The mean age was 42.3 ± 11.7 yr. There were 55 men with a mean age of 43.3 ± 11.6 yr and 6 women with a mean age of 32.8 ± 8.6 yr. The most common conditions were anxiety ($N = 29$ with a mean suspension of 276.8 d) and alcohol-related causes ($N = 10$ with a mean suspension of 493.5 d). The mean time of suspension was 385.6 ± 594.3 d. There were 20 pilots who did not return to work. The most common reasons were anxiety ($N = 8$), depression ($N = 4$), and alcohol-related causes ($N = 3$).

Ear Nose and Throat (ENT)

There was a total of 59 pilots with ENT conditions (Table VIII). The mean age was 44.1 ± 12.1 yr. There were 56 men with a mean age of 47 ± 12.1 yr and 3 women with a mean age of 34.7 ± 11.5 yr. The most common conditions were sinusitis ($N = 15$ with a mean suspension of 41.8 d) and vertigo ($N = 12$ with a mean suspension of 135.1 days). The mean time of suspension was 150.4 ± 285.9 d. One pilot did not return to work; he was diagnosed with an acoustic neuroma.

Obstetrics and Gynecology

There was a total of 30 female pilots with obstetric and gynecological conditions (Table IX). The mean age was 36.2 ± 4.9 yr. The most common condition was pregnancy ($N = 27$ with a mean suspension of 356.7 d). The mean time of suspension was 419.4 ± 382.6 d. Four pilots did not return to work; all were pregnant.

Table II. Period of Temporary Incapacitation for Pilots – Musculoskeletal.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Foot and Ankle			
Metatarsal Fracture	13	73.1±50.9	0
Ankle Fracture	12	64.9±54.4	1
Ankle Sprain	4	35.2±13.4	0
Tarsal Fracture	4	73.2±64.1	0
Achilles Tendon Tear	3	215±171.4	0
Syndesmosis Injury	3	22±16.6	0
Calcaneus Fracture	2	34.5±16.3	0
Hallux Valgus	2	219±42.4	0
Bunion Surgery	1	11	0
Foot Contusion	1	444	0
Nailbed Injury	1	40	0
Osteoarthritis Ankle	1	DNR	1
Toe Fracture	1	17	0
Unspecified Foot Injury	1	44	0
Unspecified Sprain	1	43	0
Arm and Elbow			
Proximal Radius Fracture	9	40±23	0
Forearm Fracture	3	417.5±482	0
Humerus Fracture	2	91±21.2	2
Triceps Tendon Tear	2	94±82	0
Elbow Dislocation	1	48	0
Olecranon Bursitis	1	65	0
Olecranon Fracture	1	76	0
Pain Elbow	1	3	0
Osteoarthritis Elbow	1	283	0
Hand and Wrist			
Metacarpal Fracture	16	68.1±79.8	0
Distal Radius Fracture	11	54.5±22.8	0
Carpal Bone Fracture	9	180.1±254.2	0
Phalanx Fracture	7	49.1±24.3	0
Phalanx Dislocation	4	94±86.1	0
Crush Injury Finger	3	75±36.6	0
Nail Bed Injury	3	59.6±22	0
Tenosynovitis	3	239±286.9	0
Thumb Fracture	3	20.7±37.6	0
Carpal Dislocation	2	50±14.1	0
Dupuytren Disease	2	26.5±23.3	0
Hand Fracture	2	27±29.7	0
Nonspecified Wrist Injury	2	43.5±4.9	0
Ganglion Wrist	1	24	0
Hand Ligament Tear	1	174	0
Mallett Fracture	1	41	0
Trigger Finger	1	12	0
Wrist Pain	1	21	0
Other			
Removal of Hardware	2	17	1
Osteochondrosis	1	300	0
Other Bone Disorder	1	113	1
Hip			
Osteoarthritis	5	133.8±148.1	0
Total Joint Replacement	3	108±123.1	0
Pelvic Fracture	2	1385	1
Hip Dislocation	1	49	0
Hip Pain	1	93	0
Impingement	1	960	0
Knee			
Meniscus Injury	39	56.8±65	0
Anterior Cruciate Ligament Injury	10	134±92.7	0
Pain	6	174.7±224.3	0

(Continued)

Table II. (Continued).

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Osteoarthritis	3	232.7±285.5	0
Multiligament Injury	2	81±7.1	0
Osteochondritis Dissecans	2	43	1
Effusion	1	28	0
Joint Pain	1	DNR	1
Patella Disorder	1	100	0
Patella Tendinitis	1	24	0
Posterior Cruciate Ligament	1	191	0
Prepatella Bursitis	1	104	0
Total Joint Replacement	1	33	0
Lower Extremity			
Tibial Shaft Fracture	7	137.6±164.7	0
Tibial Plateau Fracture	6	104.5±43.5	0
Femur Fracture	4	463±267.5	0
Stress Fracture	4	80.2±10.1	0
Muscle Tear	2	14.5±9.2	0
Osteoarthritis	2	82.5±2.1	0
Other Leg Fracture	2	216.5±280.7	0
Surgical Complications	1	147	0
Shoulder			
Impingement	19	121.1±246.2	0
Clavicle Fracture	13	62±64.7	0
Proximal Humerus Fracture	9	59.1±41.7	0
Rotator Cuff Tear	7	296.4±507.2	0
SLAP Lesion	7	68.7±33.3	0
Dislocation	6	121.7±167.6	0
ACJ Dislocation	4	113.5±128.3	0
Calcific Tendinitis	2	219±260.2	0
Bone Cyst Humerus	1	67	0
Other	1	92	0
Scapula Fracture	1	22	0
Spine			
Lumbar Spine Disc Prolapse	23	184.5±239.9	0
Low Back Pain	12	173.2±253.7	1
Cervical Radiculopathy	7	54.7±40.8	0
Cervical Myelopathy	6	179.8±155.8	1
Lumbar Spine Fracture	5	152.5±161.3	1
Cervical Disc Prolapse	2	54±11.3	0
Sciatica	2	22.5±9.2	0
Thoracic Spine Pain	2	705±865.5	0
Ankylosing Spondylitis	1	70	0
Cervical Disk Replacement	1	57	0
Lumbar Spine Radiculopathy	1	47	0
Sacroiliitis	1	25	0
Spondylolisthesis	1	76	0
Torticollis	1	DNR	1
Total	392	--	13

DNR: did not return.

Dermatology

There was a total of 21 pilots with dermatological conditions (Table X). The mean age was 46.3 ± 10.6 yr. There were 20 men with a mean age of 47 ± 10.5 and 1 woman who was 34 yr old. The most common conditions were melanoma ($N = 10$) and psoriasis ($N = 2$). The mean time of suspension was

Table III. Period of Temporary Incapacitation for Pilots – Medicine.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Cardiology			
Hypertension	60	101.6±222.4	1
Angina	27	217.5±349.9	2
Arrhythmia	26	170.6±189.4	1
Syncope	14	232.6±253	2
Myocardial Infarction	4	207.5±61.5	2
Atrial Septal Defect	3	451.5±453.2	1
Intracardiac Thrombosis	3	148±123.2	0
Atrial Fibrillation	2	442.5±458.9	0
Cardiac Stenting	2	516.5±338.7	0
Unstable Angina	2	641.5±649.8	0
Abnormal Cardiovascular Function	1	DNR	1
AV Malformation	1	127	0
Cardiac Valve Disorder	1	DNR	1
Ischemic Heart Disease	1	218	0
Pulmonology			
Sleep Apnea	28	173±428.3	1
Pulmonary Embolus	8	138.2±52.8	2
Covid	7	26.6±33.5	0
Upper Respiratory Infection	3	22.7±11.7	0
Asthma	2	43±7.1	0
Pneumothorax	2	45±7.1	0
Abnormalities of Breathing	1	DNR	1
Cough	1	66	0
Other Lung Disorders	1	415	0
Pleuritis	1	DNR	0
Pneumonia	1	2170	0
Shortness of Breath	1	751	0
Gastroenterology			
Crohn's Disease	5	112.4±65.1	0
Ulcerative Colitis	3	87±54.3	0
Abnormal Liver Function	2	56	1
Acute Pancreatitis	2	9±5.65	0
Gastritis	2	151±188.1	0
Irritable Bowel Syndrome	2	303.5±33.2	0
Reflux	2	159.5±166.2	0
Diarrhea	1	783	0
Nausea/Vomiting	1	DNR	1
Pyloric Stenosis	1	20	0
Neurology			
Dizziness	9	156.1±145.1	1
Migraine	3	164	2
TIA	3	271	2
Cognitive Impairment	2	259	1
Bell's Palsy	1	DNR	1
Brain Disorder Unspecified	1	548	0
Caisson's Disease	1	53	0
Dementia	1	15	0
Headaches	1	DNR	1
Multiple Sclerosis	1	281	0
Parkinson	1	517	0
Pituitary Adenoma	1	334	0
Stroke	1	DNR	1
Trigeminal Neuralgia	1	869	0
Infections			
Amoebiasis	2	16.5±4.11	0
Unclear Fever	2	36±12.7	0
Fever Unknown Origin	1	23	0
Leishmania	1	28	0

(Continued)

Table III. (Continued).

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Lyme Disease	1	803	0
Malaria	1	1354	1
Mononucleosis	1	46	0
Syphilis	1	1	0
Tuberculosis	1	245	0
Oncology			
Colon Cancer	4	580	3
Lymphoma	3	22±29.7	1
Leukemia	2	614±796.2	0
Benign Myocardial Neoplasm	1	2167	0
Esophagus Cancer	1	DNR	1
Hodgkins Lymphoma	1	DNR	1
Lung Cancer	1	DNR	1
Endocrinology/ Rheumatology			
Diabetes	26	116.9±208.9	1
Thyrotoxicosis	6	122.3±128.8	0
Iron Deficiency	5	83±134.8	0
Gout	3	47.5±26.2	1
Hypothyroidism	3	77±124.7	0
Inflammatory Arthropathy	3	257.7±289.8	0
Rheumatoid Arthritis	2	53.5±45.9	0
Enteropathic Arthropathy	1	830	0
Hashimoto's	1	13	0
Osteoporosis	1	20	0
Vitamin Deficiency	1	281	0
Nephrology			
Kidney Failure	3	191.3±145.8	0
Proteinuria	1	22	0
Other			
Thrombocytopenia	2	79±2.8	0
Deep Vein Thrombosis	1	340	0
Unspecified	1	DNR	1
Total	335	–	37

DNR: did not return.

44.9 ± 39.6 d. Three pilots did not return to work; two were diagnosed with melanoma and one pilot had other causes (not specified in the database).

DISCUSSION

The results of this database study revealed a total of 1133 temporary and 100 permanent license suspensions over a 4-yr period in the United Arab Emirates. This equates to an annual average of 283 temporary and 25 permanent license suspensions. In 2022 approximately 10,000 pilots were employed by the 4 major commercial airlines in the United Arab Emirates. This brings the annual temporary incapacitation rate to 2.8% and the permanent suspension rate to 0.25%. Evans and Radcliffe⁸ reported an annual temporary incapacitation rate of 4.3%, which is slightly higher. However, the 1.5% difference is probably not significant and within the normal margin of statistical error. It should be noted that Evans and Radcliffe⁸ also included in-flight medical events, which was not captured by our data.

Table IV. Period of Temporary Incapacitation for Pilots – Surgery.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Hernia Inguinal	53	48.4±95.4	1
Appendicitis	24	35.8±77.9	2
Cholecystitis	17	120.1±200.2	1
Hemorrhoids	12	202.7±350.2	0
Hernia Umbilical	12	56.6±95.4	0
Anal Fistula	5	86±83.9	0
Diverticulitis	4	64.5±56.4	0
Pilonidal Sinus	4	35.7±28.3	1
Post-Surgical Procedure	4	24.7±12.9	0
Rib Fracture	4	99.2±112.8	0
Thyroid Surgery	4	734.3±1119.6	1
Head Injury	3	247±189.7	0
Phlebitis	3	52.7±26.4	0
Aspiration for Complication	2	133±63.3	0
Concussion	2	28	1
Anal Cancer	1	26	0
Aortic Aneurysm	1	DNR	1
Bariatric Surgery	1	DNR	1
Breast Hypertrophy	1	8	0
Cellulitis	1	76	0
Complication from Procedure	1	13	0
Disease Intestine	1	55	0
Dog Bite	1	24	0
Hernia Diaphragmatic	1	50	0
Intussusception	1	18	0
Mandible Fracture	1	42	0
Necrotizing Fasciitis	1	1002	0
Neoplasm Aortic Body	1	204	0
Obesity	1	455	0
Occipital Condyle Fracture	1	92	0
Post GIT Surgery	1	574	0
Post Procedure	1	41	0
Hypothyroidism			
Right Quadrant Pain	1	7	0
Scalp Laceration	1	7	0
Stricture of Artery	1	DNR	1
Varicose Veins	1	4	0
Total	174	–	10

DNR: did not return.

Accurate planning and flight scheduling is important for any commercial airline and these figures will be helpful to plan and roster pilots and consider temporary license suspensions. Another important factor is cost. We have previously shown that a dedicated musculoskeletal rehabilitation unit results in significant cost savings and earlier return to work.¹⁶ When using the figures from our publication, the calculated cost for each day of temporary grounding is calculated to \$1500 per day excluding the costs for medical treatment. These figures will be helpful for financial planning.¹⁶

Musculoskeletal conditions were the most common reasons for temporary medical license suspension, with a mean period of 115 d. Of the total 392 cases, 83% were related to trauma; 152 cases were related to fractures, and 92 cases to joint dislocations and ligament injuries. Obviously, pilots have to maintain fitness and are engaged in athletic activities that possibly account for the relatively high number of

Table V. Period of Temporary Incapacitation for Pilots – Urology.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Kidney Stones	35	96.6±183.7	0
Ureter Stones	26	99.5±216.8	1
Prostate Cancer	7	69±29.7	5
Renal Colic	6	155.7±214.7	0
Bladder Disorder	3	203.7±320.8	0
Bladder Cancer	2	DNR	2
Circumcision	2	18±4.2	0
Haematuria	2	24.5±3.5	0
Prostate Hypertrophy	2	28±29.7	0
Testes Disorder	2	17±4.2	0
Kidney Cancer	1	DNR	1
Other RENAL Disorder	1	1268	0
Scrotal Varices	1	42	0
Testicular Cancer	1	DNR	1
Urethral Stricture	1	70	0
Urine Retention	1	58	0
Total	93	–	10

DNR: did not return.

traumatic injuries in this population group. The annual incidence for traumatic injuries for aircrew has previously been reported to be 73–81 per 100,000 population for women and 24–100 per 100,000 population for men.²⁴ Previously, Hohmann et al. have shown that a dedicated musculoskeletal rehabilitation unit reduced the time to return to flying by 39% from a mean of 188 to 85 d.¹⁶ Applying the findings of Hohmann et al.¹⁶ to this study, it is possible that return to flying for musculoskeletal conditions could be reduced to 70 d, resulting in significant cost savings.

The second most common reason for temporary incapacitation was medical illnesses. However, of the 335 cases, 44% were related to cardiac disease such as arrhythmias (29%), angina (24%), and hypertension (41%). The incidence of cardiac related disease is moderately higher than in the normal population. Hemingway et al. reported a 2.03 incidence for men and 1.89 in women per 100 population.¹⁵ Annual medical license examinations in pilots are a possible cause for these discrepancies, as it can be expected that quiet cardiac disease will be detected, resulting in further tests and temporary suspension. The least common reasons were dermatological conditions. Of the 21 cases, 10 were related to melanoma and 2 pilots (20%) did not return to work. This is consistent with the current published literature²⁷ and the results of this database study confirm a higher risk for pilots. A total of 59 cases were related to ENT conditions; 34% were caused by acute infections such as sinusitis, tonsillitis, and otitis. Evans and Radcliffe have reported annual incapacitation rates in pilots and noted that 6% of all medical incapacitations were associated with ENT conditions.⁸ The findings in our study were similar. In our study cohort nearly 5% of temporary suspensions were correlated to ENT conditions.

A total of 174 cases of temporary incapacitation were pertaining to surgical pathologies: 30% were related to inguinal hernia repairs, 14% to appendicitis, and 10% to cholecystitis.

Table VI. Period of Temporary Incapacitation for Pilots – Ophthalmology.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Cataract	14	136±207.5	1
Retina Disorder	14	106.6±151.3	0
Myopia	9	45.7±25.3	0
Visual Disorder	5	96±37.9	0
Keratoconus	4	86.7±19.6	0
Chorioretinopathy	3	135.7±97	0
Refractive Disorder	3	47.6±1.5	0
Conjunctivitis	2	27±26.9	0
Pterygium	2	23±5.6	0
6 th Nerve Palsy	1	509	0
Diplopia	1	122	0
Glaucoma	1	DNR	1
Irregular Eye Movement	1	9	0
Keratitis	1	74	0
Lens Implant	1	73	0
Low Vision	1	32	0
Oculomotor Nerve Injury	1	8	0
Orbital Floor Fracture	1	50	0
Panuveitis	1	49	0
Viral Infection	1	63	0
Vitreoretinal Disorder	1	69	0
Total	68	–	2

DNR: did not return.

Inguinal hernia repairs are performed at a rate of 240 episodes per 100,000 population in the United Kingdom²⁰ and the all ages incidence has been reported to be 13 per 10,000 population.²⁶ The annual incidence of appendicitis ranges between 7–10%.⁹ The incidence of cholecystitis among people with gallstones are unknown, but of patients admitted to hospital for biliary disease, 20% have acute cholecystitis.¹⁰ The findings in this study are in accordance with the published literature and do not suggest an increased prevalence in pilots.

Ninety-three pilots (7.5%) were temporarily suspended for urological conditions. The most common reasons were renal (38%) and ureter stones (28%). Evans et al. reported a slightly lower rate of 4% for genitourinary-related causes.⁸ The percentage of urolithiasis in our study cohort was 5.1%. These figures are similar to previously reported data. Hyams et al. demonstrated that 3.7–4.6% of commercial aviation pilots were diagnosed with urolithiasis.¹⁷ Masterson et al.²¹ showed that the prevalence in Navy pilots is 2.4% and Lang et al.¹⁸ analyzed data from the Global Burden of Disease database that the incidence of urolithiasis in the normal population was 1.4%. The available data suggests that pilots probably have a higher risk of genitourinary renal and ureter stones. These differences are possibly caused by the low humidity levels in the airplane cabin, resulting in hypohydration.³¹ Another contributing factor is that pilots possibly minimize fluid intake during flight to reduce the need for micturition. This may be more prevalent since the introduction of locked flightdeck doors, making it more difficult to easily access the toilet. It will be interesting whether future studies will show an increase in renal disease in pilots.

Sixty-eight pilots were temporarily suspended for ophthalmological conditions. Cataract surgery (20%) and retina

Table VII. Period of Temporary Incapacitation for Pilots – Psychiatry.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Anxiety	29	276.8±333.3	8
Alcohol	10	493.5±1108.9	3
Depression	8	832±988.9	4
Stress	6	537±528.9	1
Counseling	5	170.7±90	2
ADHD	1	DNR	1
Bipolar Disorder	1	DNR	1
Strange Behavior	1	296	0
Total	61	–	20

DNR: did not return.

disorders (20%) were the most common causes for ophthalmological conditions; a total of 68 cases were recorded. Evans and Radcliffe⁸ reported that 2% of temporary medical unfitness was associated with ophthalmological disorders. Temporary medical license suspension for ophthalmological disorders was significantly higher in our sample and reached 5.5% of all cases. The mean age of pilots in this group was 44.8 ± 13.2 yr compared to 47 yr in Evan and Radcliffe's cohort. Risk factors for cataract are age, smoking, long-term use of steroids and other medications, exposure to ultra-violet radiation, diabetes, trauma, hypertension, and radiation therapy.⁶ One could safely assume that these risk factors are similarly distributed between Evan and Radcliffe's study cohort and the study group of this research with the exception of exposure to ultra-violet radiation. In the United Kingdom the average hours of annual sunshine are 1400 compared to 3000 h in the United Arab Emirates.²⁹ The substantial difference between the United Kingdom and United Arab Emirates could possibly explain the higher incidence of cataracts.

Table VIII. Period of Temporary Incapacitation for Pilots – ENT.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Sinusitis	15	41.8±61.4	0
Vertigo	12	135.1±152.5	0
Eustachian Tube Disorder	4	10.2±3.6	0
Sinus Polyps	4	15.5±9.9	0
Sudden Hearing Loss	4	524.25±814.7	0
Vestibular Neuritis	3	533.7±325.3	0
Acoustic Neuroma	2	276	1
Hypertrophic Nasal Turbinates	2	17.5±6.4	0
Maxillary Sinusitis	2	306±490.7	0
Other Nasal Disorders	2	20±12.6	0
Cholesteatoma	1	46	0
Condition of Glottis	1	486	0
Condition of Hard Palate	1	297	0
Condition of Larynx	1	80	0
Meniere's Disease	1	564	0
Otitis Media	1	2	0
Benign Salivary Gland Tumor	1	16	0
Tonsillitis (Acute)	1	7	0
Tonsillitis (Chronic)	1	7	0
Total	59	–	1

Table IX. Period of Temporary Incapacitation for Pilots – Obstetrics and Gynecology.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Pregnancy	27	356.7±497.2	4
Endometriosis	1	14	0
Salpingitis	1	5	0
Uterus Myoma	1	9	0
Total	30		4

There were 61 pilots who were temporarily suspended for psychiatric and mental health issues. The most common causes were anxiety (47%) and alcohol disease (16%) followed by depression (13%). The mean days of suspension was the highest, with 493 d for alcohol-related disorders and 832 d for depression. In addition, the rate of permanent loss of license in this group was the highest of all subgroups, reaching 33%. Wu et al. performed an anonymous survey and showed that depressive and mental health disorders are higher when compared to the normal population, but similar to those found in other high stress occupations.³⁰

Although only 30 cases were associated with obstetric and gynecological conditions, this cohort had the longest mean days for license suspension and the second highest rate of permanent loss of license. However, of the 30 cases, 27 pilots were incapacitated because of pregnancy. It could therefore be concluded that gynecological disease does not play a role in temporary license suspensions in female pilots.

Ascertainment of medical causes for temporary license suspensions is important and highlights medical risks that require the most attention but will also aid in determining possible future changes to medical requirements.⁸ Previous studies have reported morbidity among airline pilots. Parker et al. showed that cardiac conditions accounted for 25% and musculoskeletal cases for 11% of disability.²⁵ Sykes et al. reported that of 595 pilots in New Zealand, kidney disease was observed in 3.3% and melanoma in 1.9%.²⁸ Nicholas et al. surveyed over 10,000 airline pilots for self-reported disease and established that the most common causes for disease were melanoma, motor neuron disease, and cataracts.²³ Evans and Radcliffe⁸ documented the annual incapacitation rate in pilots. In their series of over 700 patients, accidents accounted for 18%, musculoskeletal for 18%, cardiovascular events for 14%, and psychiatric diseases for 10% of the annual rates.⁸ The incapacitation rates from these studies are similar to the results from our study, showing that musculoskeletal and cardiac-related conditions account for the majority of temporary medical unfitness in commercial pilots. Pilots reporting sick or unable to return to flying duties can affect the ability of an airline to ensure a manageable roster. Typically, reserve cover and pilots on standby allow adequate coverage for these scenarios. For routine rostering and planning, the incidence of temporary medical incapacitation, knowledge about the expected mean duration of medical unfitness, and subsequent return to work is helpful. The goal of database studies is to provide information for data-driven strategies

Table X. Period of Temporary Incapacitation for Pilots – Dermatology.

	N	MEAN DAYS & SD OF LICENSE SUSPENSION	DID NOT RETURN
Melanoma	10	43±50.6	2
Psoriasis	2	45±8.5	0
Acne	1	106	0
Allergy	1	47	0
Dermatitis	1	35	0
Lipoma	1	72	0
Sebaceous Cyst	1	15	0
Squamous Cell Carcinoma	1	12	0
Vasculitis	1	76	0
Herpes Zoster	1	12	0
Other	1	DNR	1
Total	21		3

DNR: did not return.

and the data reported in this study may be valuable for airlines in planning for and including expected sickness rates for pilots. This is in agreement with Evans and Radcliffe, who noted a lack of data when studying commercial pilot incapacitation rates.⁸

This study has limitations. It is limited by the fact that it only included episodes of incapacitation resulting in temporary suspension of both medical and flying license. It is possible that pilots did not report minor injuries and illness and continued with their duty despite a compulsory reporting requirement. As such, the real rate of illness and injury may have been underreported. The period of incapacitation was defined as the time from license suspension by the regulatory authority to the time of license reinstatement. It is possible that operational and other reasons have caused delayed reporting of incapacitation and also a delay in requesting license reinstatement to the regulatory authority, causing inaccurate data. However, this is a limitation of any database study, which may vary in the degree of detail and accuracy.¹³ In contrast, database studies are considered to be representative of the population of interest and serve as an inexpensive source of reliable data.¹³ The established criteria for return to work and reinstatement of the return to flight privileges was based on the regulations of the local regulatory body¹² and other regulatory authorities^{7,12} may have different functional criteria. This limits the external validity of this study. It is possible that the ICD-10 categorization of pathology by the treating physician was incorrect, over- or underinflating certain diagnoses. In addition, it is also possible that we have incorrectly categorized reported ICD-10 codes.

In conclusion, this database study showed that musculoskeletal conditions are the most common reasons for temporary loss of medical license, followed by medical and surgical conditions. The least common reason was dermatological conditions. The longest period of incapacitation was associated with psychiatric conditions followed by medical and ENT conditions. The annual calculated temporary incapacitation rate was 2.8% and the permanent suspension rate was 0.25%. The results of this study may help in understanding the reasons for temporary pilot incapacitations related to medical conditions and possibly aid with rostering and return to work planning.

ACKNOWLEDGMENTS

Financial Disclosure Statement: The authors have no competing interests to declare.

Authors and Affiliations: Erik Hohmann, FRCS (Tr&Orth), Ph.D., M.D., Burjeel Hospital for Advanced Surgery, Dubai, United Arab Emirates, and School of Medicine, University of Pretoria, Pretoria, South Africa; and Reino Pieterse, M.B.Ch.B., M.Sc. (Sports Medicine), Clinical Lead, Emirates Group Rehabilitation Unit, Dubai, United Arab Emirates.

REFERENCES

- Band P, Deschamps M, Fang R, Le N, Gallagher RP. Long term disability rates in a cohort of Air Canada pilots. *Aviat Space Environ Med.* 1998; 69(12):1137–1140.
- Burgard SA, Lin KY. Bad jobs, bad health? How work and working conditions contribute to health disparities. *Am Behav Sci.* 2013; 57(8):1105–1127.
- Butler GC, Nicholas JS. Health among airline pilots. *Air Line Pilot.* 2001; 2001(March):16.
- Caplan A, Batra P. The ethics of using de-identified medical data for research without informed consent. *Voices in Bioethics*; 2019 Nov. 14. [Accessed 9 Jan. 2023]. Available from <https://journals.library.columbia.edu/index.php/bioethics/article/view/5917>.
- Caruso CC. Negative impacts of shiftwork and long work hours. *Rehabil Nurs.* 2014; 39(1):16–25.
- Chang JR, Koo E, Agron E, Hallak J, Clemons T, et al. Risk factors associated with incident cataracts and cataract surgery in the Age-related Eye Disease Study (AREDS): AREDS report number 32. *Ophthalmology.* 2011; 118(11):2113–2119.
- EASA. Aircrew & medical: medical. [Accessed 9 Jan. 2023]. Available from <https://www.easa.europa.eu/en/domains/aircrew-and-medical/medical#regulations>.
- Evans S, Radcliffe SA. The annual incapacitation rate of commercial pilots. *Aviat Space Environ Med.* 2012; 83(1):42–49.
- Ferris M, Quan S, Kaplan BS, Molodecky N, Ball CG, Chernoff GW. The global incidence of appendicitis: a systematic review of population-based studies. *Ann Surg.* 2017; 266(2):237–241.
- Fialkowski E, Halpin V, Whinney RR. Acute cholecystitis. *BMJ Clin Evid*; 2008; 2008:0411.
- GCAA. NPA 10-2014 medical standards. [Accessed 9 Jan. 2023]. Available from [https://www.gcaa.gov.ae/en/epublication/EPublications/Notice%20of%20Proposed%20Amendment%20\(NPA\)/NPA%2010-2014%20MEDICAL%20STANDARDS.pdf#search=proposed%20amendment%20medical%20standards](https://www.gcaa.gov.ae/en/epublication/EPublications/Notice%20of%20Proposed%20Amendment%20(NPA)/NPA%2010-2014%20MEDICAL%20STANDARDS.pdf#search=proposed%20amendment%20medical%20standards).
- Guide for aviation medical examiners. [Accessed 9 Jan. 2022]. Available from https://www.faa.gov/ame_guide.
- Hashimoto RE, Brodt ED, Skelly AC, Dettori JR. Administrative database studies: goldmine or goose chase? *Evid Based Spine Care J.* 2014; 5(2):74–76.
- Hege A, Lemke MK, Apostolopoulos Y, Sönmez S. Occupational health disparities among U.S. long-haul trucker drivers: the influence of work organization and sleep on cardiovascular and metabolic disease risk. *PLoS One.* 2018; 13(11):e0207322.
- Hemingway H, McCallum A, Shipley M, Manderbacka K, Martikainen P, Keskimäki O. Incidence and prognostic implications of stable angina pectoris women and men. *JAMA.* 2006; 295(12):1404–1411.
- Hohmann E, Pieterse RJ. Pilots after shoulder surgery and rehabilitation in a dedicated musculoskeletal rehabilitation unit of a major airline returned to work earlier when compared to standard rehabilitation by external providers. *Arthrosc Sports Med Rehabil.* 2022; 4(1):e1–e7.
- Hyams ES, Nelms D, Silberman WS, Feng Z, Matlaga BR. The incidence of urolithiasis among commercial airline pilots. *J Urol.* 2011; 186(3):914–916.
- Lang J, Narendrula A, El-Zawahry A, Sindhwani P. Global trends in incidence and burden of urolithiasis from 1990 to 2019: an analysis of global burden of disease study data. *Eur Urol Open Sci.* 2022; 35:37–46.
- Ludvigsson JF, Haberg SE, Knudsen GP, Lafolie P, Zoega H, et al. Ethical aspects of registry-based research in Nordic countries. *Clin Epidemiol.* 2015; 7:491–508.
- Maisonneuve JJ, Yeates D, Goldacre MJ. Trends in operation rates for inguinal hernia over five decades in England: database study. *Hernia.* 2015; 19(5):713–718.
- Masterson JH, Phillips CJ, Crum-Cianflone NF, Krause RJ, Sur RL, L'Esperance JO. A 10-year retrospective review of nephrolithiasis in the Navy and Navy pilots. *J Urol.* 2017; 198(2):394–400.
- National Health and Nutrition Examination Survey 2011–2012. [Accessed 5 January 2022]. Available from: <https://wwwn.cdc.gov/nchs/nhanes/ContinuousNhanes/Default.aspx?BeginYear=2011>.
- Nicholas JS, Butler GC, Lackland DT, Tessier GS, Mohr LC Jr, Hoel DG. Health among airline pilots. *Aviat Space Environ Med.* 2001; 72(9):821–826.
- Ong T, Sahota O, Marshall L. Epidemiology of appendicular skeletal fractures: a cross-sectional analysis of data from the Nottingham Fracture Liaison Service. *J Orthop Sci.* 2015; 20(3):517–521.
- Parker PE, Stepp RJ, Snyder QC. Morbidity among airline pilots: the AMAS experience. *Aviation Medicine Advisory Service. Aviat Space Environ Med.* 2001; 72(9):816–820.
- Primates P, Goldacre MJ. Inguinal hernia repair: incidence of elective and emergency surgery, readmission and mortality. *Int J Epidemiol.* 1996; 25(4):835–839.
- Sanlorenzo M, Wehner MR, Linos E, Kornak J, Kainz W, et al. the risk of melanoma in airline pilots and cabin crew: a meta-analysis. *JAMA Dermatol.* 2015; 151(1):51–58.
- Sykes AJ, Larsen PD, Griffiths RF, Aldington S. A study of airline pilot morbidity. *Aviat Space Environ Med.* 2012; 83(10):1001–1005.
- Wikipedia. United Kingdom's climate. [Accessed 9 Jan. 2023]. Available from https://en.wikipedia.org/wiki/Climate_of_the_United_Kingdom.
- Dubai's climate. [Accessed 9 Jan. 2023]. Available from https://en.wikipedia.org/wiki/Climate_of_Dubai.
- Wu AC, Donnelly-McLay D, Weisskopf MG, McNeely E, Betancourt TS, Allen JG. Airplane pilot mental health and suicidal thoughts: a cross-sectional descriptive study via anonymous web-based survey. *Environ Health.* 2016; 15(1):121. Erratum in *Environ Health.* 2017; 16(1):129.
- Zubac D, Stella AB, Morrison SA. Up in the air: evidence of dehydration risk and long-haul flight on athletic performance. *Nutrients.* 2020; 12(9):2574.